



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

LANE

MEDICAL



LIBRARY

LEVI COOPER LANE FUND

REASONS FOR PRESCRIBING

Fairchild's Pepsin.

It is the most active, gives the greatest value for the price asked.

One grain will digest more albumen than will a grain of any other pepsin made. We do not care by whose test, what the proportions of albumen, water, and acid, only let the conditions be the same for all.

It is positively guaranteed to be permanent. It will not get soft or sticky, or deteriorate in value, because it is not a peptone. It is free from all the well-known characteristics of peptone. It is not made by a patented or pirated process.

If you prescribe pepsin in scale or powder, and it becomes sticky, you may be sure that it is not Fairchild's. Investigate the matter, place the blame where it belongs.

There are pepsins "made to sell" under the title of scales—products which are inferior and dissimilar to Fairchild's pepsin in scales. They are peptones made by a patented process, and sold under cover of a title first and properly used only by Fairchilds to describe their original true pepsin in the form of scales.

The druggist is told, "They are just as good as Fairchild's. If Fairchild's is not specified, put in mine." But claims and pretensions, however colossal, cannot make peptones permanent.

There is no soluble or peptone pepsin equal in activity to Fairchild's, none which will keep in all weathers and all conditions as Fairchild's will.

FAIRCHILD BROS. & FOSTER,

New York.

DRS. BOURNEVILLE AND BRICON'S

MANUAL OF

HYPODERMIC MEDICATION.

BY

G. ARCHIE STOCKWELL, M. D., F. Z. S.
(Member of New Sydenham Society, London.)



1890.
GEORGE S. DAVIS.
DETROIT, MICH.

MP



Copyrighted by

GEORGE S. DAVIS.

1890.

W. A. A. J.

TABLE OF CONTENTS.

INTRODUCTION.

	PAGE.
History, Value and Methods; Solutions; Injections; Instruments and Care of.....	1-13

MANUAL OF SUBCUTANEOUS INJECTIONS.

Acidum Benzoicum; Acidum Carbolicum; Acidum Hydrocyanicum; Acidum Osmicum; Acidum Phos- phoricum; Acidum Tannicum; Aconitum; Agar- acinum; Alcoholum; Aloes; Ammoniaë Liquor; Amyl Nitrite; Aqua; Arbutinum; Argenti Nitras; Arsenicum; Aspidosperminum and Qubrachinum; Auri Chloridum.....	15-35
Belladonna; Brominum.....	35-38
Caffeina; Calabarina, (Physostigmina, Eserina); Camphora; Cannabis Indicaë; Cedrinum; Chloral Hydras and Chloral Crotonis; Chloroformum; Cin- chona Salts; Coniina; Convallaria Majalis; Cotoin- um and Para-Cototoinum; Curara.....	38-61
Digitalis; Duboisina.....	62-63
Emetina; Ergota; Ether; Eucalyptol.....	64-66
Ferrum.....	66-67
Gelsemium.....	67
Hydrargyrum; Hydrastinina; Hyoscyamus.....	68-81
Ichthyolum; Iodoformum.....	82-83
Jaborandum.....	83-85

VI.

	PAGE.
Moschi Tinctura; Muscarina.....	85-86
Nicotina; Nitro-Glycerinum; Nux-Vomica.....	86-65
Oleandrina; Oleum Amygdalæ and Oleum Olivæ; Opium.....	86-94
Paraldehydum; Pereirina; Picrotoxinum; Potassii Bromidum and Iodidum; Potassii Permanganas, 102-105	
Quinolinum; Quinonum or Hydroquinonum.....	106-113
Resorcinum.....	
Scillianum and Scillipicrinum; Sodium Salts; Solan- ina; Sparteina; Stramonium.....	114-126
Tayuya; Terebinthinæ Oleum.....	121
Urethan.....	121
Valdivinum; Vaselineum; Veratriinum.....	122-125
Zinc Salts.....	125
INDEX.....	127

PREFACE.

The *Manuel des Injections Sous-cutanées* of Drs. Bourneville and Bricon has achieved such notable success on the Continent and in Great Britain, that no apology is required in introducing a modification thereof to cis-Atlantic practitioners, especially as there are no works in the English language devoted to this subject at present available. With the exception of Bartholow's *Manual*, out of print since 1884, and Eulenburg's essay in Ziemssen's *Handbook of Therapeutics*, nothing has been published in this country within recent years.

This work possesses the advantage of being very complete, well up to date, and above all, of pointing out certain disadvantages that are liable to accrue from practice of this branch of therapeutics.

It is with sincere pleasure I acknowledge my indebtedness to Dr. Andrew S. Currie, of Moorlands, Sidney, Gloucestershire, England, whose English publication has made my own work simple and easy. *Moreover, the additions and emendations of this gentlemen are in many instances the most valuable paragraphs in the book.*

The work has been almost entirely re-written, as it seemed advisable to omit a considerable portion of original matter that was either irrelevant or repetitious, and also the reference notes, which are largely superfluous now that nearly every public library possesses files of the *Index Medicus*, that afford more complete reference. The formulæ too, where it seemed advisable and no way modified the context, have oftentimes been changed to meet the requirements of American practitioners, who for the most part employ decimal measures; further, 5 and multiples of 5, are more easily calculated and employed than the multiples of such figures as 4, 6, and 8.

Trusting the volume will prove as valuable and interesting to the reader as it has to the editor and revisor, I remain, sincerely,

G. ARCHIE STOCKWELL.

Detroit, Mich., August 1st, 1890.

INTRODUCTION.

HISTORY, VALUE AND METHODS.

The introduction of a needle into the body as a therapeutic measure, has been in vogue for centuries in the Orient, and was introduced into Great Britain and Europe in 1670, though it met with little favor until 1810, when Berloz revamped the procedure. Intravenous injections generally obtained in the 17th century, but whether a *hollow needle* bearing medicaments was ever employed for remedial purposes, is questionable. Endermic medication, however, was taught by Lambert; "Inoculation of Remedies," by Lafargue, in the early part of the present century; and Valleix attempted the introduction of medicines into the cellular tissue by acupuncture.

For the strictly hypodermic method, however, aside from the physiological experiments of Magendie and others, we are indebted to Doctor Alexander Wood, of Edinburgh, Scotland, who employed morphia solutions subcutaneously for the relief of neuralgia as far back as 1843, though such procedure attracted little attention until about 1855. Priority has been claimed for Langenbeck, Rynd, Kurzak, J. Taylor, and others, but has never been satisfactorily substantiated. Rynd wrote of it in 1845, but there is little doubt his information was derived from Dr. Wood. No small portion of our present knowledge, moreover, is due to Mr. Charles Hunter and Doctor An-

stie, the former of whom became the sponsor of the method, deriving the name from *υπω* (under) and *δερμα* (the skin).

Since 1855-56 when the hypodermic method was introduced into America by Doctor Fordyce Barker, and an impetus given thereto by Ruppaner who employed the methods of Valleix, the recorded experience of a large number of trustworthy observers throughout the civilized world has raised to the rank it at present justly holds. Its popularity also is in some measure due to the Commission appointed by the Royal Medico-Chirurgical Society of Great Britain, in 1867, who reported as follows:

1. That as a general rule only *clear neutral solutions* should be injected, for such solutions rarely produce local irritation.

2. That whether the drugs be injected under the skin or administered by the mouth or rectum, their chief physiological and therapeutical effects are the same *in kind*, though varying in *degree*.

3. That symptoms are observed to follow the subcutaneous injection of some drugs, which are absent when the same are administered by other methods; and on the other hand certain unpleasant symptoms which are apt to follow the introduction of the drugs by mouth and rectum, are not usually experienced when such are injected under the skin.

4. That, as a general rule,—to which, however, there may be exceptions,—clear neutral solutions of drugs introduced subcutaneously are more intense in their effect than when introduced by the rectum or mouth.

5. That no difference has been observed in the effects of a drug subcutaneously injected, whether it be introduced near to, or at a distance from, the part affected.*

6. That the advantages to be derived from this method of introducing drugs, are: *a.* Rapidity of action: *b.* Intensity of effect: *c.* Economy of material: *d.* Certainty of action: *e.* Facility of introduction in certain cases: *f.* With some drugs the avoidance of unpleasant symptoms.

7. This plan, therefore, is especially adapted to those occasions when very rapid and decided effects are required from drugs that are operative in small doses.

SOLUTIONS.

With regard to the safety of hypodermic procedures, there are no dangers providing due caution is exercised, backed by thorough knowledge of the physiological manifestations of the materials employed, and of the chemical compounds that may

* Messrs. Eulenberg and Davis, deriving their ideas from Lambert, hold absorption is more rapid in certain situations than others. For instance: Most rapid when introduced in the temples or cheeks; next, in the epigastrium; third, in the anterior portion of the thorax; fourth, in inner aspect of arm and thigh; fifth, in the nape; sixth, external aspect of arm and thigh; seventh, in forearm; and eighth, *nil* in the foot and back.

The absurdity of this is self-obvious. Since all injections obtain their effect by absorption into the circulation, it is evident the differences that may arise from locality are practically inappreciable.—ED.]

be induced in the solutions by keeping, by the effects of exposure to air, by sunlight, etc. Again, it must be remembered that the physiological activity of nearly all drugs, more especially narcotics and those that possess direct action upon the great nerve centers, is from three to five times more active when administered hypodermically, than when given by the mouth; likewise, that the liquid injected must not be markedly acid nor deeply alkaline, nor in any way obviously irritant to tissue.

It is a mistake, however, to suppose that it is always absolutely indispensable that a solution should be strictly neutral; it is merely desirable it should be *as nearly so as possible*. In fact, we are often obliged to have recourse to acid solutions, which, unfortunately, as a rule, are more or less painful; but it may be observed that suffering is not generally proportionate to the acidity of the solutions—thus, quinine in tartaric acid is more painful than in sulphuric or hydrochloric acid, although the acidity of both the latter is more intense. Again, the mere fact a solution is neutral, is not evidence also that it is non-irritating.

The degree of concentration is also to be considered. In regard to substances of very active properties, this is of minor importance; but we are precluded from employing most substances hypodermically, because the bulk demanded for a dose is too great; because the solubility is too slight to permit of being

used in sufficient doses; or because the remedy in the necessary degree of concentration is highly irritating.

In selecting a solvent, the question of *result* is imperative. As far as possible, all injections are best made fresh, and, when permissible, camphor-, or chloroform-waters prove the best vehicles, providing it is not a question of alterative injection. Solutions in equal proportions of glycerin and water are frequently well borne by the tissues, but not always desirable owing to the caustic action of the former; and other solvents and vehicles may also be satisfactorily employed, such as distilled water, cherry-laurel water, ether, alcohol, oils, etc.; and when a preparation is to be kept for any length of time, it is often advantageous to add chloral, carbolic acid, boric acid, or some other material that will secure its preservation.

Accidents may arise from deposits or fungi; crystals may appear in a too concentrated solution, having either existed from the first, or subsequently developed; or the medicine itself may be at fault, especially if it is held in suspension instead of solution—as in the case of calomel—or if it is but moderately capable of absorption. Finally, as far as possible, it is best to inject fluids at a temperature near that of the body, or about 98° F.

INJECTIONS.

The best mode of making the injection is that laid down by Mr. Charles Hunter, viz: "To pick up

a fold of loose skin and push the needle through until the point works loosely in the subdermic cellular space;" then to inject steadily, but as quickly as possible, placing the finger for a moment on the opening after the withdrawal in order to prevent the oozing of the final drop that tends to follow the needle. There is no need for a plaster or anything of the sort over the puncture.

An adept will hold the syringe barrel between the fingers with the head of the piston pressed against the ball of the palm, and by an adroit movement insert the needle and solution by one instantaneous movement as it were, the whole procedure occupying less than two seconds of time. This is the writer's method, and is so rapid that even children find no cause for complaint, especially as a platinum cannula as fine, almost, as a cambric needle is employed.

Formerly it was taught to inject slowly, with two or three pauses of a second or two, but this is unnecessary (and withal oftentimes most painful) unless the amount employed is considerably in excess of fifteen or twenty minims.

If the operator be expert, and his needles suitable, there need be no pain worthy of mention; it is only by carelessness that a needle is badly inserted,—transfixes a nerve or vein, or enters connective tissue. Slight hæmorrhage, a drop or two, is of no moment; and the same in the main holds true when a small blood-vessel is transfixed, if the fluid is free from turbidity and

non-irritating; again injections of air, as a rule, are of little consequence, and, beside, need never occur if the operator understands his business, and properly fills his syringe.

It is, perhaps, unnecessary to remark that cachectic, purpuric, diabetic, and alcoholic individuals are predisposed to sundry accidents, and that in infectious diseases the syringe should never be employed except under strict antiseptic precautions. Also, that injections should never be made on the nose, ears, eye-lids, different parts of the neck, scrotum, axilla, groin, fingers, etc., nor in œdematous regions or parts which are the seat of stasis, inflammation, or extravasation.

INSTRUMENTS.

Of the many syringes recommended, it is unnecessary to speak in detail, since all are very nearly of the same pattern. Three, however, deserve especial mention.

First, and to my mind, the best, is Leiter's, (Fig. 1.), with platina needles of exquisite sharpness and fineness, the barrel (D) cut from a solid block of glass, the bore beveled at either end, to accommodate the hard-rubber beveled fittings (*a*, *b*). It can always be kept tight; there are no washers and no cement to loosen and thus open the joints; the packing can always be spread and re-introduced without impairing its shape or utility; and there are no

these tablets is, division of material with the greatest possible uniformity and exactness; further they keep indefinitely in any climate, and the materials of which they are composed are absolutely non-irritating.



FIG. 4.—Full size.

It may be well to remark, *en passant*, that the syringes designed to receive tablets in the heel of the needle, and to dissolve the medicament by forcing fluid through it into the cellular tissue of the patient, are utterly unreliable when employed in this way. The tablets oftentimes are not dissolved, or fine particles are forced into the tissues, and perhaps also left in the needle to its infinite damage. The tablet is best dissolved in a spoon, or within the barrel of the syringe itself, before employing as an injection. With P., D. & Co.'s syringe, the plunger may be removed, the tablet inserted, the required amount of water added, and when solution is complete the plunger replaced,

provided the guard cap (B—Fig 5.) that protects the needle-end of the instrument is not removed.

Figure 5 represents a plated metal case of flattened elliptical shape suitable for carrying in vest or card pocket, to replace the morocco case of figure 4.

The syringe of Messrs. P., D. & Co. provides, also, a special arrangement against leakage in case the plunger should become dry from disuse. Attached to the piston rod on the upper side of the plunger (C), is a small nut (A) that, on the rod being drawn up into the cap, fits there into a socket (F), when by a turn of the piston (D) the packing is expanded. Other advantages are: The ease with which the needle is cleaned—on the insertion of the wire into the needle cap it is immediately guided by bevelled sides to the opening in the needle proper; the reinforcement of the needle by a sheath embracing the upper half, strengthening in the part most likely to give way under pressure; and the fastening of the needles in the case by screw threads whereby their points are preserved from injury. Another advantage derived from this fastening, which is a left-handed screw, is the barrel can be screwed to the needle, and the latter removed without handling, aiding in preserving the needle aseptic. The syringe is furnished either with or without finger rests.

In filling care should be taken to exclude all air bubbles. In Leiter's instrument this is best done by drawing back the plunger before attaching the bottle



FIG. 5—Full Size.

(at *B*) to the barrel (at *a*), then forcing the air into the bottle, withdrawing the piston again, when the syringe, if held perpendicularly with the bottle above, is properly filled; the needle can now replace the bottle. Some introduce the needle into the cellular tissue before fitting to the syringe, but this is clumsy; it should be adjusted to the barrel beforehand, after having duly ascertained its perfect cleanliness, and that the instrument is in perfect order otherwise.

The operation performed, it remains only to cleanse the needle, and if necessary, disinfect it. It is always desirable to expel the last drop of fluid, and to rinse inside and out with clear water, followed by alcohol; and it should not be laid aside without inserting a fine wire, or, better yet, a pig's bristle. It is also a good plan, in order to always insure perfect asepsis, to keep the needles when not in use, immersed in a phial of alcohol. Needles that have become obstructed, may be cleansed by holding for an instant in the flame of an alcohol lamp, when the foreign substance is usually quickly driven off. If a wire has rusted in a needle, it should be dipped in oil before being held in the flame; and to remove the rust from the interior, oil may be passed through, heating again, then rinsing with alcohol.

EDITOR.

MANUAL OF SUBCUTANEOUS INJECTIONS.

ACIDUM BENZOICUM—BENZOIC ACID.

An anti-fermentive and anti-putrescent; slightly soluble in cold water (1 to 200), more so in boiling water (1 in 4), and in glycerin (10 per cent.); very soluble in alcohol, ether, and turpentine.

Is eliminated by the perspiration; by the saliva also as succinic acid; by the kidneys as hippuric acid; and by the lungs when the dose exceeds 15 to 30 grains. It gives rise to no remarkable phenomena, and is employed chiefly as an expectorant and excitant. The strongest solution employed by Rohde caused great pain, but never gave rise to abscesses or ecchymoses.

℞ Benzoic acid, 18 grains.

Alcohol, q. s. ft., 120 minims.

Dose, 5 to 15 minims ($\frac{3}{4}$ to $1\frac{1}{2}$ grs.). Both solution and syringe should be warmed to 98° F.

This solution may cause indurations that persist for several days, and sometimes induces puffiness and redness of skin.

Has been employed in pneumonia, typhus, uræmia, collapse, etc. It has also been recommended in the treatment of chronic bronchitis, cystitis, and gout (Currie).

BENZOATE OF SODA, which is soluble in two parts of water, appears to be possessed of more active properties than benzoic acid, but, as far as known, has not yet been employed hypodermically.

ACIDUM CARBOLICUM—CARBOLIC ACID.

PHENOL. PHENIC ACID.

An anti-putrescent and anti-fermentative. Soluble in all proportions in alcohol, ether, and glycerin, and in 50 parts of water.

Absorption is rapid, even through the unbroken cuticle, and elimination is correspondingly so, through the lungs, kidneys, and skin. The urine, without being necessarily heavily charged with the acid, often assumes a greyish-green hue, and sometimes there is albuminuria, especially after large doses.

From seven to thirty grains induces general excitement, followed by stupor, vertigo, *tinnitus aurium*, deafness, languor, sweats, diminution of pulse and temperature, colic, and diarrhœa. Poisonous doses result in vertigo, giddiness, tingling of fingers, intoxication, delirium, stupor, analgesia, anæsthesia, muscular debility, intermittent heart followed by arrest in diastole, collapse, coma, death—rarely convulsions as in lower animals.

Antidotes:—Saccharated lime (Husemann); sulphates (Baumann); inhalations of oxygen, bleeding, transfusion (Ferrand).

℞ Carbolic acid, 1 grain.

Distilled water to make 140 minims.

Dose, 10 to 20 minims, which produces a sensation of heat or burning, usually disappearing after a half hour; in large doses may procure small ecchymoses without abscess.

℞ Carbolic acid, 1 grain.

Glycerin to make 70 minims.

Five minims represents $\frac{1}{14}$ grain.

This has been used in traumatic erysipelas, which it arrests inside of five days according to Bœckel, who employed 15 minims in five or six injections in the healthy tissue at a distance of one centimetre from the diseased patch. He repeats the second time during the same day, pursuing the treatment until fever subsides and there is an arrest of erysipelatous inflammation.

[Recently Baccelli (*Semaine Medical*, 1888, p. 422, and 1890, p. xxiv.) has lauded the use of carbolic acid in subcutaneous injections in neuralgias in general, and particularly in the sciatic, supra-orbital, and intercostal forms. He also reports a case of tetanus cured by this means. A one per cent. solution was employed, making the injections over the seat of pain, repeating every hour till relief was obtained. He believes the efficacy of the acid is due to the sedative action which it exerts on the excitability of the nervous centres. But little pain or smarting follows the injections.—Ed.]

Carbolic acid has also been employed in intermittents (Jessier, Déclat, Heuter); prurigo (Rezek,

Fleischmann); diphtheria (Trotz); phthisis and tuberculosis (Schnitzner, Eliacopulos); crural neuralgia, pleuro-pneumonia (Hagen, Kunze); diabetes, charbon, malignant œdema, anthrax (Raimbert, Méplain, Trélat); nævus (Bradley, Stockwell); acute articular rheumatism (Bergonzini, Senator, Kunze); Ludwig's angina.

CARBOLATE OF AMMONIA has been employed hypodermically in typhoid.

ACIDUM HYDROCYANICUM.

HYDROCYANIC (PRUSSIC) ACID.

Hydrocyanic acid is soluble in all proportions in alcohol and water, and has a somewhat irritant action similar to that of morphine. Dr. Bartholow has employed that of the U. S. P. in four drop doses, but considers it too large, even a dangerous dose; in fact it is not prudent to exceed *two* drops, which, however may frequently be repeated, owing to its rapid elimination.

R Acid hydrocyanic dilute, U. S. P., 20 minims.
Camphor water, 40 minims.

Dose, from 5 to 12 minims (Ed.).

CYANIDE OF POTASSIUM, less potent than hydrocyanic acid, is readily soluble in water, in alcohol, and in glycerin (32 per cent.); and is prescribed hypodermically in doses of from $\frac{1}{4}$ to $\frac{3}{4}$ of a grain cautiously increased to $2\frac{1}{2}$ grains; but even in such feeble doses

as one per cent., it gives rise to severe pain, though without suppuration.

R Potassium cyanide, 1 grain.

Camphor water, 180 minims.

Dose, 5 to 12 minims (ED.).

CHERRY-LAUREL WATER is extensively used as a solvent for morphine, the mixture forming a clear solution [that, however permanent, is objectionable owing to the tendency to oxydize.—ED.] The cherry-laurel water of the American and English Pharmacopœias contains exactly one per cent. of hydrocyanic acid, and is non-irritating except when it has become acid through oxydation by contact with air. Dose, 5 to 10 minims, cautiously increased.

[The ESSENTIAL OIL OF BITTER ALMONDS of commerce is a mixture of volatile bitter almond oil (*hydride of benzoyl*) and hydrocyanic acid, with small quantities of benzoic acid, benzoin, and benzamide. It is highly poisonous, being in general four times as strong as officinal hydrocyanic acid, for which it has occasionally been employed as a substitute; but at best it is a dangerous and uncertain remedy. Dose, $\frac{1}{30}$ to $\frac{1}{8}$ minim in alcohol and glycerin; the smallest amount is likely to produce urticaria and other unpleasant symptoms.—ED.]

The effects of all three of these drugs are practically the same: Bitter taste, feeling of heat and burning, followed by numbness of the tongue, pharynx, and stomach; also salivation, increased number of respira-

Fleischmann); diphtheria (T. S. S. painful—elevation of
culosis (Schnitzner, Eliaco... and spasmodic phe-
pleuro-pneumonia (Hagen, K... they induce nausea, vom-
malignant œdema, anthrax... difficult respiration—either
lat); nævus (Bradley, Stock... paralyze respiration and
matism (Bergonzini, Sena... changes—dilatation of
gina. ... pressure and temperature,

CARBOLATE OF AMMONIA. Pulse, general tonic and
hypodermically in typhoid. ... motor paralysis, coma,

ACIDUM HYDROCYANICUM. Therapeutically in mental dis-
HYDROCYANIC ACID. ... mania and eclampsia (Mac-

Hydrocyanic acid. ... anacholia (*Ibid*, Bartholow,
alcohol and water, and ... pectoris, gastralgia, func-
similar to that of morphia. Brinton, Stille, Bartholow);
ployed that of the U. S. ... uria, lichen tropicus, eczema,
considers it too large. Vest, Fuller); nervous cough,
fact it is not prudent. Whooping cough, dyspepsia of
however may frequently be used. (Watson).—Ed.]
rapid elimination. ... asserts atropine is a physio-

B. Acid hydrocyanic. The experiments of Bartholow do
Camphor. Inhalation of ammonia is an
Dose, from 5 to 10 grains. Very little use by the stomach; it
is absorbed hypodermically, or by intra-

CYANIDE OF POTASSIUM. Cyanide of potassium is a
cyanic acid, is re-
in glycerin (32 grains). OSMICUM—OSMIC ACID.

mically in dose. ... employed by interstitial injection in the
increased to 24 grains; and by Billroth and others in
... with varying results.

Twenty minims may be used at one time, which represents $\frac{1}{14}$ grain.

ALCOHOLUM—ALCOHOL. SPTS. VINI RECTI.

For hypodermic use is best employed at a temperature of 90° F.; at a higher temperature there is danger of burning the tissues, and even at this, if too near the extremities, there is the same risk. At a lower temperature it tends to diffuse, and its local effects are proportionately less intense; but then one is administering a stimulating injection with general action. The doses must be calculated according to the results desired, hence no definite figure can be fixed. It produces neither abscess or sloughs—merely a smarting sensation which speedily subsides—and is largely used as a diffusible stimulant. Flood made more than 200 injections of whisky and coffee in four hours in a case of opium poisoning, with complete success, though at the same time he injected *tincture of belladonna* in 10 and 20 minim doses, which may have contributed to the result. Edward Warren (1867) and Ainsworth injected brandy in collapse, and Breisky and Figuero had good results from the same in acute anæmia due to puerperal hæmorrhage or traumatism.

ALCOHOL AMMONIO-ANISETUM.—Ammoniated alcohol with anise, is prepared by Zuelzer, of Berlin, according to the following formula:

Essence of anise, 10 minims.

Alcohol at 85°, 40 minims.

Solution of ammonia, 50 minims.

Antidotes.—Richardson has used hypodermic injections, and Wood, and, however, intravenous injections, of ammonia; “even when the cardiac contractions are affected, or have actually ceased, it is possible to establish them by the artificial stimulus of ammonia.”—(Maury.)

NAPELLINE.—[An amorphous alkaloid, soluble in water, ether, alcohol, etc. The tincture it possesses over the fluid preparation of aconitine, are more than problematical.

AGARACINUM—AGARACIN.

Has been extolled by Seifert, Piering, and by whom it was used to check excessive perspiration, and who claim also that it diminishes the frequency of cough in phthisis, without inducing either vomiting or diarrhoea; sleep is more tranquil and uniform. As the maximum activity is only attained 2 to 3 hours after administration, it is necessary to administer it at various times according to the nature of the case. Seifert discourages the use of the tincture together, since given by the mouth it induces vomiting, and table diarrhoea, and employed hypodermically it produces an intense burning sensation.

- R Agaracin, 1 grain.
Absolute alcohol, 80 minims.
Glycerin, up to 200 minims.

VATEI
dermic
sects ar
stroke (W
hæmoni
eneral col.

arat., 1 drachm
rachms.
as required.

ng by *tincture*
inims of ammori
minutes. This wa
of ten minutes.
poisoning had dis
at one point of in
kin.

AMMONIA has also be
s of from 1 to 5 grain
alts are characterized
properties, that they are
treatment of collapse

L. NITRITE.

a powerful sedative and a
s temperature by ozonizing
king oxydation, and relaxes
y reducing arterial pressure.

a rule it is much more effective by inhalation than by the mouth or subdermically. Indeed, the results obtained by hypodermic use are uncertain, though Dr. Barnes claims to have secured effects precisely parallel to those occurring to administration by inhalation. Drs. Bourneville and Bricon assert "from 16 to 80 minims have been injected in the dog, and 90 minims in man, in one and a half hours," but this statement is open to grave doubt. Personal experience has taught that ten minims hypodermically may prove unsafe, even if given in divided doses, in the time indicated, as it tends to paralysis of respiration.

I have found the following of great benefit oftentimes in dysmenorrhœa,—and Mary Putnam Jacobi has obtained equally satisfactory results from amyl by inhalation:

R	Amyl nitrite.....	10 parts.
	Alcohol.....	30 "
	Glycerin.....	60 "

Dose 8 to 15 minims (in lumbago and sciatica, deeply injected.)

S. Weir Mitchell, Jones, and Maraghano have obtained good results in epilepsy; it often proves effective in sea-sickness, but like all remedies for this malady is not inevitable in overcoming. It is invaluable in whooping-cough, if given *per os*, just before the onslaught of the paroxysm, hence might be tried subdermically. Its antidotes are chloroform, and cocaine.—Ed.]

ANTIMONIUM TARTARATUM.—TARTAR
EMETIC. TARTRATE OF ANTIMONY.

Soluble in water 1 in 15; in glycerin $5\frac{1}{2}$ per cent.; insoluble in alcohol.

It may truly be said that tartrate of antimony is hardly ever indicated except as an emetic.

R Tartrate of antimony, 1 grain.
Tartrate of morphine, $\frac{1}{8}$ grain.
Distilled water, up to 40 minims.

Dose, 10 minims ($\frac{1}{4}$ grain).

AQUA—WATER.

Has been repeatedly substituted with success for morphine in localized painful affections, in doses of from 15 to 35 minims. It causes some little sharp pain, but never gives rise to ill-effects. ["Dr. Fergus of Glasgow," says Dr. Currie "declares that in many cases acute pain can speedily be relieved by the injection of *hot* water. The syringe should first be warmed thoroughly, and the temperature of the water raised to as high a degree as can be borne without causing actual pain. In one case of colic this method afforded relief from suffering very speedily, but in cases where it was used for the relief of tabetic pains the effect was at the most equivocal."—Ed.]

ARBUTINUM.—ARBUTIN.

A glucoside from *Arctostaphylos Uva-ursi*, soluble in water, yielding hydroquinon by decomposition in

the system. Our knowledge regarding this drug is meagre and unsatisfactory, but it is generally deemed a valuable diuretic, relieving catarrhal conditions of the bladder, and even renal hæmorrhages. In cases of purulent inflammation of the bladder and kidney, it exercises a direct antiseptic influence, lessening the formation of pus. Dose, subcutaneously, 4 grains.

ARGENTI NITRAS.—NITRATE OF SILVER.

The pain which follows hypodermic injections of silver nitrate are often very sharp, although of short duration. "They are irritating to the degree of almost necessarily producing an abscess at the point of puncture; nevertheless, the silver limits, in a manner, its own action by the zone of coagulation which it produces. There is no kind of inflammation more limited—in the centre, a slough and a sero-purulent collection which isolates and eliminates it; at the circumference a ring of induration, often almost of cartilaginous consistence" (Luton). The artificial abscess allows of the escape of serous or purulent fluid, carrying with it a slough of cellular tissue, leaving a cicatrix.

℞ Crystallized nitrate of silver, 10 grains.

Distilled water to make 120 minims.

Dose, 5 to 20 minims.

A solution of 1 in 5 almost always determines slough and abscess; and rarely the former is not separated but remains encysted in the living tissue like a

foreign body. With a solution of 1 in 20, it is to be observed that the irritation decreases the nearer the injection is made to the trunk; thus, when made at the hip there is no certainty of an abscess resulting. M. Luton does not think that there is any advantage gained by using the weaker solutions. Without referring to its use for surgical purposes, nitrate of silver has been employed hypodermically in neuralgia (Luton, Ruppaner, Bertin and Michaliski), rheumatism, gout (Ruppaner), chronic arthritis, tabes dorsalis (Frommhold).

ARSENIIUM.—ARSENIC.

R *Fowler's solution*, 2 drachms.

Camphor water, 3 drachms.

Dose 5 to 20 minims.

Eulenburg claimed to have obtained good results in the various form of tremor (paralysis agitans, &c.); Kobner, in lichen ruber, and in a case of diffuse sarcoma of the skin (child of 8 years); Lesser, in lupus; Soltzmann, in chorea, etc.,—no particular symptom was observed, however, save that patients occasionally complained of nausea. [Bartholow recommends *liq. sodii arseniatis* in preference to Fowler's solution, for hypodermic purposes, as it is less irritating in its local effects, less apt to produce arsenical poisoning, and possesses greater osmotic effect. The injections are made on alternate days, and in different situations so as to minimize the local

effects. Personally, in most instances where an *alterative* effect is desired, and in the majority of skin diseases, Donovan's solution, (*liq. arsenii et hydrargyri* Ed.) is preferred, and it may be employed in the same way as Fowler's solution.—ED.]

Arsenic is recommended in: Chorea (Radcliff Hammond, Smith); cholera—algid stage (von Graëfe); various forms of tremor (Eulenburg); paralysis agitans (Beauveville); psoriasis (Lewin, Lipp, Tichomilew), chronic eczema (Lipp); neuralgia, epilepsy, psychoses, etc (Radcliffe, Tebaldi).

In small doses it causes a sensation of heat in the epigastrium and stomach, increase of appetite, increase of gastric functions, diminution of urea and carbonic acid, lowering of temperature and pulse. In large doses, feeling of constriction of throat, thirst, epigastrium, nausea, vomiting, colic, diarrhœa, fever, tachycardia, insomnia. In poisonous doses, diarrhœa with rice-water stools that sometimes may be bloody, prostration, pallor of the face, debility, irregular menstruation, scanty and suppression of urine, dyspnoea, vertigo, unconsciousness, delirium, convulsions, coma.

Acute poisoning. Cutaneous eruptions (eczema, erythema, urticaria, &c.); conjunctivitis, extreme anorexia, vomiting, diarrhœa, inflammation and ulceration of the oral and laryngeal mucous membranes; paralysis of the extensors; dropsy, cachexia, fatty degeneration of most of the organs. Elimination is by

the bile, urine, and perspiration, and its duration is most variable. The physiological effects of arsenic further than this are but little known.

ASPIDOSPERMINUM AND QUEBRACHINUM.

Aspidospermin the active principle of *Aspidosperma quebracho*, is soluble slightly in water, freely so in alcohol, ether, and essential and fixed oils, not at all in glycerin. The phenomena observed after administration are, decreased frequency of the pulse and respirations (Berthold, Picot, Berger, Laquer). Continued use induces headache, vertigo, mental hebetude, and profuse salivation (Laquer, Berger). Quebrachin, another active principle, is analogous in action to curare. According to M. Closson, paralysis of the limbs in those which survive in poisoning by aspidospermin, would be due to an action of the poison on the nervous reflex centres in the spinal cord, the excitability of which is lost before the peripheral nerves, motor and sensory, and the muscles, have been attacked by the poison. But with sufficient doses of aspidospermin, the peripheral nerves are first paralyzed, then the muscles lose their sensibility to stimuli to an equal extent.

The *sulphate* and *hydrochlorate* are amorphous readily soluble salts; their solutions are intensely bitter, and are deemed useful in dyspnoea of all kinds (Bourneville).

R Sulphate of aspidospermin, 5 grains.

Camphor water, up to 160 minims.

Dose, 8 to 16 minims *plus*.

[Dr. Currie justly remarks: "The statement that quebracho and its preparations may be used in all forms of dyspnœa, must be accepted with caution. It has however, been used in dyspnœa dependant on various forms of cardiac disease, *e. g.*, hypertrophy, aortic incompetence, etc. From the experiments of Harnack and Hoffmann, it appears that the greatest therapeutic value of quebracho bark consists in its property of reducing the irritability of the respiratory centre. It will prove beneficial in cases of dyspnœa *not* depending on impeded aëration or insufficient oxydation of the blood in the lungs, and that is not a mere compensatory symptom; namely, in dyspnœa in consequence of disorders of circulation and of diseases of the heart. In other cases, in which the dyspnœa constitutes a purely compensatory manifestation, any attempt at lessening the irritability of the respiratory centres may, under certain circumstances, be fraught with danger, although this remedy may even then occasionally assist in relieving this distressing symptom." Aspidospermin is an emetic, and more energetic than apomorphine, but does not act in the same way on the brain and respiratory centres.—ED.]

AURI CHLORIDUM—CHLORIDE OF GOLD.

R Chloride of gold, 1 grain.

Distilled water, 560 minims.

Dose, 16 minims ($\frac{1}{3}$ grain).

[Recommended in the treatment of certain syphilitic symptoms; and has been successfully used by M. Moricourt in hysterical anæsthesia in a solution of 1 part in 500 and 1000.—ED.]

BELLADONNA.

℞ Neutral sulphate ATROPINE, 1 grain.
Camphor water, 2 ounces + 160 minims.

Sixteen minims of this solution contains $\frac{1}{40}$ of a grain of atropine sulphate.* It is always wise to begin with a small dose, say $\frac{1}{40}$, and to rarely exceed $\frac{1}{8}$ of a grain. "In the following formula, morphine combined with sulphate of atropine, permits us to obtain all the good effects of these two drugs, especially in the treatment of pain and insomnia, and to increase the dose with safety" (Dujardin-Beaumetz):

Hydrochlorate of morphine, 1 grain.
Sulphate of atropine, $\frac{1}{40}$ grain.
Cherry laurel water, up to 224 minims.

Sixteen minims contain $\frac{1}{4}$ of a grain of morphine and $\frac{1}{40}$ of a grain of atropine. [This combination, has long been extolled, but is alike unchemical, unphysiological, and unpharmaceutical. The effects claimed are indefinite, and depend solely upon the idiosyncrasies of the patient that permit one or the other drug to assert itself to the exclusion of the

*It will be well to remember that very little *atropine* is to be had in market, the alkaloid usually sold under the name being *hyoscyamine*—ED.]

B Sulphate
Campho.
Dose, 8 to 16

[Dr. Currie
that quebracho
all forms of d
caution. It has
dependant on va
hypertrophy, ac
periments of H
the greatest r
sists in its prom
respiratory co
of dyspnœa
insufficient ox
that is not a
dyspnœa in
and of disea
the dyspnœa
festation,
the respir
stances, l
may even
tressing
more
same w

AU

are those of atro-

have been used
complaint that
however, has not
with a saturated
or when the nor-
was used.—ED.]
indicated in
intestinal incontinence
intestinal obstruc-
of phthisis,
atropine is of
anti-sialagogue,
attributes to it
than ergot when

including its
produce dryness of
of the pupils; in
deglutition, facial

when there is dry-
generally, with
employed in such cases, it
"Currie". This
belladonna, however.
is necessary
light-complex-

anæsthesia, blindness, cephalalgia, vertigo, delirium, hallucinations, nausea, slowing followed by acceleration of pulse, scarlatiniform redness of the skin, œdema; *in poisonous doses*, aphonia, muscular spasms, collapse, coldness of the surface, and finally death with arrest of the heart in diastole. Symptoms of poisoning appear in a few minutes. Elimination is effected rapidly by the kidneys.

No accident is likely to occur, however, provided the solution used is perfectly pure, contains neither acid nor alcohol, and is freshly prepared.

Antagonisms.—Calabar bean, physostigmine, and pilocarpine; as to muscarine, antagonism has not been conclusively proved.

HOMATROPINE, according to Fronmüller, does not include any of the poisonous elements of atropine; its local mydriatic action is quick and decided (2 per cent. solution), and the pupil rapidly returns to the normal condition. It diminishes tubercular sweats.

Hydrobromate of homatropine, P., D. & Co., 1 gr.,
Distilled water, up to 80 minims.

Sixteen minims= $\frac{1}{2}$ of a grain.

Neither mydriasis or narcotic phenomena are induced; cough, expectoration and insomnia are relieved.

Dr. William Murrell has found these injections less efficacious than those of atropine. Homatropine has been used, among other remedies, in cases of poisoning by pilocarpine (Fronmüller).

BROMINUM.—BROMINE.

Soluble in 40 parts of water, in alcohol, ether, carbon bisulphide, chloroform, glycerin. Used by Goldsmith hypodermically in hospital gangrene in one drop doses, in the immediate neighborhood of the wound, when, after forty-eight hours the latter lost its characteristic appearance.

[Dr. Carvallo, U. S. A., informed me that, witnessing some of the experiments, he was shocked at the pain induced, and that while bromine changed the character of the wound, it also induced ulceration equally objectionable and terrible.—ED.]

CAFFEINA—CAFFEINE AND SALTS.

Soluble in 80 parts of water and in 50 of alcohol, but is very unstable. Most of the solutions produce a sense of burning due to the addition of acid or alcohol.

℞ Caffeine, $1\frac{1}{2}$ grains.
Alcohol, 6 minims.
Camphor water, up to 96 minims.

Dose, 8 to 20 minims, *℥ss*.

℞ Caffeine, 20 grains.
Salicylate of sodium, $17\frac{1}{2}$ grains.
Camphor water, to 1 drachm.

Dose, 1 to 6 minims; contains 1 grain in 3 minims.

Particularly recommended for alcoholic and morphine intoxication, also for hemicrania (Currie).

Garrison and M. Pallen have injected liquid extract of coffee in cases of acute morphinism with success.

Therapeutically, caffeine, or a liquid extract of coffee (Parke, Davis & Co.'s *fluid extract*) may be used in neuralgia (Eulenburg, Anstie, Pletzer); migraine (Lorent); insomnia in chronic alcoholism without delirium (Anstie); melancholia, hysteria, cholera (Oser); strangulated hernia (Guzman); heart disease where digitalis is contra-indicated (Riegel)—in the latter case it is well to begin with small doses, and if necessary increase them rapidly; its greatest effect is obtained by small doses frequently repeated, and it acts more rapidly than digitalis without producing cumulative effects (Currie). Further, caffeine has been strongly recommended as a diuretic—daily doses of 7 to 12 grains of the citrate and hydrochlorate, induced such abundant diuresis that general œdema disappeared in a few days, and being non-cumulative, it may be steadily administered for months (Gubler, contradicted by Riegel, who employed in 8 cases of pleurisy and 4 of nephritis, and found inferior to other diuretics). Also, it has been recommended in the treatment of asthma by Thorowgood. Its action on the temperature is the reverse of quinine (Currie). In small doses, ($1\frac{1}{2}$ grains), there is increase of functional activity of the brain without any kind of trouble; increase of force of circulation and of vascular tension.—Doses of 6 to 16 grains, by subcutaneous injection, induce moderate slow-

ing of cardiac contractions, and increased fulness and tension of the pulse (Riegel).—In large doses, increase or diminution of pulse rate, lowering of blood pressure, elevation of temperature (Binz), headache, tremor of the hands, nausea, vomiting, drowsiness, tinnitus aurium, *muscæ volitantes*, priapism, frequent desire to micturate, hallucinations, vertigo, acceleration then diminution of the respirations; increases also reflex excitability. Poisonous doses in animals induce general paralysis, arrest of the heart in diastole, death.

Elimination takes place by the urine without modification, and by the bile (Strauch). The *sulphate*, *hydrochlorate*, and *hydrobromate*, decompose in water. The *citrate* is merely a mixture of caffeine and citric acid (Jehl).

CALABARINA. PHYSOSTIGMINA.—CALABARINE. PHYSOSTIGMINE. ESERINE.

Physostigmine, or eserine, is slightly soluble in acidulated water, and very soluble in alcohol, ether, and chloroform. Most of the salts are soluble in water. The solution of *sulphate* of eserine on exposure to air and light assumes a violet-red color, more or less intense, the transformation into red eserine being much less active in the case of eserine than of its sulphate. The *salicylate* of physostigmine is soluble in 150 parts of water and in 12 parts of alcohol. The solutions of this salt are yellow, but

become red under the influence of light. Calabarine, another alkaloid (Harnack and Witkowski), is rarely employed medicinally.

℞ *Extract of Calabar bean*, 2 grains.

Distilled water up to 80 minims.

Dose, 10 minims ($\frac{1}{4}$ gr.) every two hours which may be increased to $\frac{3}{4}$ gr.

℞ *Physostigmine*, 1 grain.

Alcohol, q. s.

Distilled water to make 128 minims.

Dose, 16 minims, $\frac{1}{8}$ grain (Erlenmeyer).

℞ *Eserine*, 1 to 5 grains.

Distilled water (acidulated?) to 112 minims.

Dose, 4 minims, $\frac{1}{8}$ to $\frac{1}{4}$ grain.

[Bourneville and Bricon assert eserine and its sulphate may be injected in doses of from $\frac{1}{32}$ to $\frac{1}{16}$ grain, but these are perhaps too large, and apt to produce either a slight degree of paralysis of the diaphragm, or some embarrassment of this muscle. "The hypodermic use of eserine sulphate produces a very sharp pain which lasts from one to six hours" (Suarez y Cruz).

In spite of all claims and the multitude of formulæ, I have been unable to secure a satisfactory solution except as follows:

℞ *Extract of Calabar bean*, 2 grains.

Camphor water, 60 minims.

Make fresh as required, since no solution will keep any length of time. Five minims represents $\frac{1}{4}$

grain. Messrs. Parke, Davis & Co., prepare hypodermic tablets of *sulphate of physostigmine*, each representing $\frac{1}{60}$ grain, which are most satisfactory.—
ED.]

As in the case of coniine, the results given by various investigators are very contradictory regarding physostigmine *et al.*, and this is doubtless due to the varying composition of the preparations employed, and also to the varying proportion of alkaloids contained by the Calabar bean. Nevertheless, the preparations of Calabar bean act alike on the eye, salivary glands, intestines, heart and respiration; their action on the spinal cord depends on the proportion of physostigmine or Calabarine. They are employed in: Ptosis (Schelske), enuresis (Fronmüller), poisoning by strychnine (Newman), tetanus (Watson, Monti, Rottrock, Franzolini, Duffy, Suarez y Cruz, Th. Anger, Reulos), chorea (Harley, Ogle, Bouchut, Cadet de Gassicourt and Suarez y Cruz), trismus neonatorum (Eschenburg).

Bromide of eserine, soluble in water (especially warm water) and in alcohol, insoluble in ether, chloroform, fixed and volatile oils, slightly soluble in glycerin, possesses all the properties of eserine and its salts; is perfectly neutral, and produces no local irritation. Its solution is more stable than that of the other salts of eserine (Duquesnel). There is no evidence that it has ever been employed hypodermically.

The physiological phenomena induced by Calabarine and physostigmine are: Pallor, then redness of the skin, abdominal pains, vomiting, embarrassed respiration, vertigo, feeling of extreme languor, myosis, salivation, sweats, slowing of the pulse (Fraser—toxic phenomena observed in *himself*); watery and muco-sanguinolent diarrhœa; slowing, and finally cessation of respiration; complete paralysis and collapse (Evans). Death as by asphyxia. Paralyzing effect on the central nervous system; stimulating then paralyzing effect on the peripheral nerves. Dose of $\frac{1}{140}$ to $\frac{1}{70}$ of a grain is sufficient to induce poisonous symptoms in man (Harnack). In a case of epilepsy with idiocy, physostigmine administered in doses of $\frac{1}{140}$ of a grain three days in succession, increased alarmingly the number of fits. Elimination is effected by the saliva and bile.

The mutual antagonism between physostigmine and atropine is admitted by most authors, but is denied by Messrs. Frohlich and Rossbach. Some have also asserted a reciprocal antagonism between physostigmine and strychnine.

CAMPFORA.—CAMPFOR.

A feeble antiseptic soluble in alcohol, ether, the volatile and fixed oils, and in acetic acid. Hypodermically, it induces sharp pains without consecutive local accidents.

rhagia, uterine hæmorrhage, threatened abortion, inertia or atony of uterus prolonging labor—(Chur-
chill, McGuire, McClintock, C. M. Stockwell, G.
A. Stockwell), cancer of uterus (Grailey Hewett)
palpitation of heart, eczema (Christison), cholera
(Willeman, O'Shaughnessy), ulcer of stomach (Bim-
ton), hysteria (Reynolds), dropsical affections and
Bright's disease (Bryan). In fact it is a drug of
wondrous scope and utility, too little known or
understood.—Ed.] Dr. Strange, of Worcester, ob-
tained good results in the early stage of mania
with delusions, as did Frommüller from cannabis
tannate in nervous insomnia (Currie), and in one
case of chronic miliary tuberculosis, where the
prolonged use of morphine had been productive of
no result. Eulenburg obtained by hypodermic ex-
hibition, calmer sleep and perceptible diminution of
cough and pain.

℞ [*Normal liquid cannabis Indica*, 10 minims.
Glycerin,
Camphor water, ℥ss 25 minims.

Dose, 5 to 15 minims as required; no unpleasant symptoms
develop.—Ed.]

In moderate doses is followed by a sense of
warmth, tingling and pricking, weight in head, tinnitus
aurium, oppression, uneasiness, constriction of the
throat, dryness of the mouth, nausea, vomiting, diar-
rhœa (?)—these are not always observed, however.
In *larger* doses it is a powerful nerve excitant; there

is very great activity of the intellectual faculties, mirthful and sometimes furious delirium, sense of physical and moral well-being, and of lightness; illusions and hallucinations, chiefly visual; increased acuteness of hearing [increased sexual desire—satyriasis—Ed.]; the movements of locomotion are spasmodic or incoordinated; sleep sometimes disturbed by night-mare. In *poisonous* doses, deeper coma allied to that of narcotism, anæsthesia and analgesia; rarely there is a manifestation of cataleptic phenomena, lassitude, cephalalgia. The opinions of authors are contradictory as to the effect of cannabis Indica on the circulation and temperature. The bronchial, urinary, and sweat secretions are increased.

Chronic poisoning.—Physical and mental enfeeblement, listlessness and melancholia, very marked emaciation, stiffness, and sometimes tremblings of the limbs and slowness of movements.

CEDRINUM.—CEDRIN.

An amorphous product of the seeds of *Simaba cedron*, readily soluble in water. In a dose of $\frac{1}{16}$ grain hypodermically administered, it caused vertigo in an adult. [Its tonic nature is indisputable, and it is to a moderate extent a febrifuge; it is frequently successful in relieving ague and intermittent fever if persisted in, along with the use of hepatic stimulants. Its claimed antidotal properties in poisoning by venomous reptiles, etc., are mere pretension and superstition.—Ed.]

ANAL. HYDRATE. CROTON CHLORAL.

have been employed hypodermically, but in second administration. The former, however, is added in small proportions to solutions to preserve them. Dr. Currie has observed that the unpleasant effects might be minimized, if they are alleviated, by practicing deep intra-muscular injections in the gluteal region, but this is not the writer's experience.—ED.]

CHLOROFORMUM—CHLOROFORM.

employed hypodermically by Hunter, who observed on account of the acute pain inflammation to which it gave rise. It is chiefly in neuralgia of the *trigeminal*, *maxillary*, *Weir Mitchell*, and *Mattison*. In the injection of half a drachm, noted in four minutes, puffiness round the injection for two hours: the neuralgic pain. Collins injected successfully 30 grains of old-standing sciatica. M. of his patients, after injection were not painful (50 min. in middle of thigh), complete relief. In another, a painful induration of long standing which after a course of treatment, he observed, but it must be observed to make a large number of

injections. [Let me remark that in long standing sciaticas in the *aged*, serious abscesses may result.—ED.]

CINCHONA SALTS.

QUINIDINE, which is isomeric with quinine, possesses very nearly the same physiological properties, but neither it nor its salts are suitable for hypodermic purposes.

QUININE.—English quinine is very sparingly soluble in water (1-576) but readily dissolves in alcohol; American is scarcely at all soluble in either water or alcohol; both are readily dissolved by dilute acids and chloroform. It is an anti-putrescent, anti-fermentative (Binz) anti-zymotic, [and oxytoxic—ED.], sialagogue (slightly), and diminishes tactile sensibility; it dulls the sense of hearing, causes tinnitus aurium, auditory hallucinations, vertigo, sense of weight of head, confusion of ideas, carotid pulsations, cephalalgia, quinine intoxication, ataxic gait, apathy, drowsiness, general prostration, dilatation of the pupils, and nausea: In larger doses, vomiting, delirium, deafness, occasional blindness, and aphasia: In *poisonous* doses, collapse, convulsions, and death.

There is increased volume of the red corpuscles due to increased supply of oxygen, and also disappearance of amœboid movements of the white corpuscles, the number of which is diminished. In *moderate* doses there is first increase, and then slowing of the cardiac pulsations, due to lowered blood pressure;

finally, there is arrest of respiration, followed by paralysis of the heart. In healthy persons the effect on the temperature is either *nil* or trifling. [It is a powerful antipyretic, and properly employed, vastly superior to any of the newer drugs of this class.—ED.] Elimination is effected rapidly and chiefly by the kidneys, the secretion of urine becoming very abundant, while the secretion of sweat is arrested. In some rare instances various eruptions (eczematous and scarlatiniform), and hæmaturia, have been observed after the administration of quinine. [Some cases of tetanus have been ascribed to the sub-cutaneous use of quinine, but the result undoubtedly was chargeable to the septic character of the instrument employed rather than to the medicament it bore. *Muriate* of quinia is less apt to produce cinchonism than the other salts, is the most stable, and the proportion of contained quinia moreover, greater; it is the most suitable of all the salts of quinine for any purpose whatever. A larger portion of the *sulphate* on the market is little else than *sulphate of cinchonidia* (65 per cent.)—ED.].

As a general rule, great circumspection is required in using hypodermic injections of quinine and its salts, for, in many instances, it produces, besides pain, persistent indurations, followed by suppuration and even gangrenous sloughs. [Especially is this true of the sulphate.—ED]. It is advisable to inject no more than 15 minims at one point, and to expel the fluid very gently. In cases of urgency, such as pernicious fevers,

we ought not to be deterred from using hypodermic injections by the risk of local accidents. Glycerin is often the best vehicle; it dissolves one part in six, by weight, by means of heat, but in the cold it dissolves only 3 per cent. M. Denis, who has employed solution of the sulphate in glycerin, has discontinued this form, as he could only obtain weak solutions ($\frac{1}{4}$ grain to 8 minims) even with the aid of heat.

R Muriate of quinine, 32 grains.
Dilute hydrochloric acid, 15 minims.
Distilled water, up to 160 minims.

Five minims, one grain.—ED.

This solution reddens litmus paper and preserves its limpidity without any precipitate for several months. Dr. Lente, of N. Y., uses the following:

R Sulphate of quinine, $10\frac{1}{2}$ grains.
Dilute sulphuric acid, 18 minims.
Carbolic acid, 1 grain.
Water, up to 112 minims.

Sixteen minims = $1\frac{1}{2}$ grains.

Out of 300 injections there were only two instances of abscess, and one of slough. One is naturally astonished at these results, with a fluid containing such a large proportion of acid, and in an amount which is more than enough.

The *carbolate* of quinine [which, however, is far from being a true chemical salt—ED.] has been used by M. Maestro Perez in the treatment of cholera:

Carbolate of quinine, 5 grains.

Alcohol, up to 16 minims.

Three injections are given one after the other, in the first stage of cholera. [They should be intra-muscular instead of subcellular.—Ed.]

The *formiate* of quinine is fairly soluble, and has been used by Messrs. Namias, Lévi, and Calza. The *quinat*, *tannate*, *valerianate*, *hydrobromate*, *sulpho-vinate*, *bi-sulphate*, and *lactate* of quinine have been tried, but are not of sufficient importance for more than mention here.

The *carbamide-hydrochlorate* of quinine (a combination of hydrochlorate of quinine with urea), which is soluble in its own weight of water, has been injected in a 50 per cent. solution by M. Jaffé, 16 minims being equivalent to $5\frac{1}{2}$ grains of the quinine salt. The local reaction is, as a general rule, trifling, neither redness nor puffiness being observable. In doses of 16 grains, hummings of the ears are frequently observed, but only in women. In intermittent fevers its action is always uniform, and the paroxysms arrested after three or four injections. In typhus the temperature falls considerably after the injection of two or three syringesful. [Recently has come to my notice a claim made by several physicians that this salt produced frightful abscesses, sloughing, and septic poisoning. While it is more than probable that the hypodermic syringe was at fault, it is still well to observe extreme caution in its use.—Ed.]

Quinine has been extensively used subcutaneously by many physicians in the treatment of intermittent and remittent fevers, and in tubercular diseases; also in septicæmia, mania, neuralgia, cholera, infantile cholera, typhoid fever, acute and chronic articular rheumatism, sunstroke, convulsions, and pyæmia. However, the hypodermic mode of administration should only be practiced in cases of urgency, or when there is difficulty of administering by the mouth or rectum.

QUINODINA.—*Chinoidine*, or *Amorphous Quinine*.—The *hydrochlorate* is soluble in equal parts of water, and is readily absorbed, but may produce abscess. Heybom employed in hypertrophy of the spleen.

COCAINA—COCAINE AND ITS SALTS.

Extracted from coca leaves, this alkaloid occurs in the form of white colorless needles, sparingly soluble, of a faintly bitter taste. The *hydrochlorate* is of a *creamy* white color; crystallizes in rhomboidal needles, and dissolves readily in water. It produces local anæsthesia. Other physiological effects are: Dilatation of the pupils by paralysis of the filaments or the great sympathetic (Berthold); elevation of the temperature in small doses, lowering after large and poisonous doses (minimum or maximum of temperature variation one-half to one degree after injection) (Nègre); increase in pulse volume; increase in respiratory movements followed by irregularity; lowering, followed quickly by increase, of blood

pressure (Laborde); vertigo after doses of $\frac{1}{4}$ to $1\frac{1}{2}$ grains (von Anrep). In large doses is produced hyper-excitability, epileptiform convulsions (from its action on the brain and spinal cord), general analgesia and mydriasis (Moreno y Maiz, Rondeau and Glay, Laborde, Vulpian). According to M. Berthold, there is similarity of action between cocaine and atropine; injection into the jugular vein of an animal of $\frac{1}{4}$ to $\frac{1}{2}$ of a grain of hydrochlorate of cocaine, produces considerable fall of blood pressure without a previous rise. [Amyl nitrite is both antagonistic and antidotal to cocaine.—ED.]

Solutions (4 per cent) of cocaine have repeatedly been made in and around tumors for their painless removal. Others have availed themselves of the same means in opening whitlows and abscesses.

℞ Hydrochlorate of cocaine (P., D. & Co.'s), 3 grs.,
Camphor water, 120 minims.

Dose 10 minims = $\frac{1}{4}$ grain.

Cocaine has been employed in the treatment of melancholia, dysphagia, migraine, odontalgia, hemi-crania, etc. "Prof. Mosler obtained excellent results from the use of *salicylate* of cocaine in the treatment of asthma, employing $\frac{1}{4}$ of a grain of a 5 per cent. solution (Currie). [Personal experience does not altogether corroborate this. ED.]

COLCHICINA—COLCHICINE.

℞ Colchicine, 1 grain.
Distilled water, up to 560 minims.

Sixteen minims for one dose = $\frac{1}{8}$ grain.

Injection causes sharp burning pain, and in a little more than 30 per cent. of cases signs of local inflammation also.

[This drug, and its derivative (*Colchicum seminis*), have long enjoyed an undeserved reputation in the treatment of both chronic and acute articular rheumatism, and also in neuralgia, including sciatica.—ED.]

CONDURANGO.

[The injection of a *decoction* of condurango has yielded encouraging results in some few cases of gastric and œsophageal cancer, but the probabilities are the favorable manifestations were purely matters of coincidence. Bourneville and Bricon regard it as especially valuable in gastric catarrh, but here it is best administered *per orem* in the form of *wine*.—ED.]

CONIINA. CONIINE (CONICINE).

Insoluble in water, it is freely so in alcohol and ether.

℞ Coniine, 1 grain.

Proof spirit, 120 minims.

Camphor water, 180 minims.

Dose, 5 to 10 minims = $\frac{1}{60}$ to $\frac{1}{30}$ grain,

The dose of *hydrobromate* should not exceed $\frac{1}{30}$ to $\frac{1}{6}$ grain, and possesses no local irritant action, which may occur with coniine. The latter is caustic as well as irritant.

℞ Hydrobromate of coniine, 1 grain.

Camphor water, 20 minims.

Dose, 1 to 3 minims.

Coniine and its salts have been employed as narcotics (Busch); praised in the treatment of convulsive cough, dyspnœa, laryngismus stridulus, various forms of asthma, spasm—tetanus, eclampsia, epilepsy, hysteria—and neuralgia (Tuloup, Tyriakan, Dujardin-Beaumetz); also in pulmonary emphysema, angina pectoris (Erlenmeyer, Lorent), pneumonia and pleurisy (Stewart), fever (Wertheim), blepharospasm (Eulenburg), whooping cough, bronchitis, spasmodic laryngitis (Méga), *tic douloureux* (Chaussier, Duméril), chorea (Welch, Harley), acute mania (Brown).

The physiological effects are most uncertain and variable: Dejection; sadness; general tremors; dyspnœa; rapid pulse; increase, then diminution, and finally loss, of reflex irritability; slowing of the pulse and respiration; analgesia; paralyzing action on motor nerves, which lose their excitability. The vagus nerve, which is the first to suffer, is also the first to recover. Pure coniine has no physiological action on the motor and sensory nerves. Respiration is most rapidly affected, the heart the last to die; the organs of sense, the digestive and calorific functions, are little affected (Tuloup); affections of sight; collapse; and death by asphyxia. The hydrochlorate and *hydrobromate* of coniine are stable salts, and their action is similar to that of coniine, but more energetic.

CONVALLARIA MAJALIS.—LILY OF THE VALLEY.

Convallaria majalis, recommended by various authors as a cardiac remedy, has been used hypodermically in a few cases by Dr. Andrew Smith. "Cases Illustrating the Action of Convallaria on the Heart" (Archives of Medicine, viii, p. 293).

[Farquharson (Woodbury) remarks it is a cardiac tonic and diuretic acting usefully in failure of compensation in mitral stenosis and regurgitation. Napheys claims it as an excellent substitute for digitalis, and adds, "a *fluid extract* such as is prepared by Parke, Davis & Co., may be administered (by the mouth) in fifteen or twenty drop doses every three hours until its effect is produced." Dr. Constantine Paul asserts it is a true heart tonic, but must be used in very large doses; Dr. S. T. Bruen, that in valvular diseases it is most useful in cases of mitral obstruction, but loses its effectiveness in proportion as fatty degeneration is present. "In advanced valvular heart disease, or cardiac failure, in the catarrhal nephritic forms of Bright's disease, convallaria, has not, however, proven as useful as digitalis." Margilano believes it increases the amount of urine.

That it is of great service in cases where palpitation and dyspnœa, rather than deficient cardiac systole, are the prominent features, is certain; it is especially valuable in phthisis or asthma in which palpitation and dyspnœa are prominent symptoms, before cardiac failure from advanced fatty change

ensues. Moreover, its effect is oftentimes brilliant in “purely functional heart disorder, especially palpitation with irregular cardiac action dependent upon general debility, and in cases of anæmia or hysteria. It is sometimes an efficient remedy in the irregular action of the heart caused by tobacco” (Napheys). The active principle is CONVALLAMARIN, a glucoside soluble in water.

℞ Convallamarin, 1 grain.

Glycerin, 40 minims.

Camphor water, 140 minims.

Dose, 5 drops = $\frac{1}{8}$ grain, cautiously increased, and not to exceed $\frac{1}{4}$.

Its physiological action is that it diminishes the number of pulsations while augmenting the amplitude of the contractions.

Albuminuria prevents or lessens its diuretic action. No cumulative effects have been observed.—
ED.]

COTOINUM. PARACOTOINUM—

COTOIN. PARACOTIN.

[Cotin and para-cotin are to all practical purposes identical, save that the latter is one-fifth weaker in therapeutic value. Both are bitter crystals, soluble in boiling water, alcohol, ether, and chloroform, slightly so in cold water; anti-putrescent and anti-zymotic, and invaluable in the treatment of intestinal fluxes, except in some cases of ulceration, or of cirrhosis of the primæ viæ.

℞ Cotoin (or paracotoin, 6 grains), 4 grains.

Acetic ether to make 15 minims.

Dose, 15 minims repeatedly hourly (or even every 15 or 20 minutes), or as required, the injection made deeply.—ED.]

Insoluble in the gastric juice, cotoin is freely so in the intestinal fluids, and acts by dilating the abdominal vessels, and lowering the temperature (Albertoni, Burkent).

[It is a stimulant alterative and stomachic, increasing the appetite, and specific in diarrhœas in that it modifies and restores the functional energy of epithelium, and stimulates pancreatic and biliary secretion.—ED.] Frömmüller asserts it to be antisudorific, and Albertoni adds also it is anti-sialagogue, and both recommend in ptyalism and cholera, while M. M. Burkart and Jobst, who employed in the latter disease, suggest its combination with chloral [probably because coto has no effect on peristalsis, while chloral—or analgesin—are directly anæsthetic to the intestines.—ED.]

Elimination is effected chiefly by the kidneys, and there is marked diminution of indican in the urine. [I have employed paracotoin in equal parts of glycerin and water.—ED.]

CURARE OR OURARI (*Strychnos toxifera*.) CURARINA.—OURARINE. WORARA.

Ourari is soluble in water, in alcohol and water, insoluble in ether. *Ourarine* is deliquescent and readily soluble in alcohol and water. In doses of from

$\frac{1}{4}$ to $\frac{5}{8}$ of a grain, ourari, or curare, causes congestion of the brain; violent but transient headache; feeling of fatigue, apathy; increased salivary, lachrymal, sudoriparous and urinary secretions; glycosuria; pulse of greater force and frequency; more rapid respiration, and increase of temperature. In doses of $1\frac{1}{2}$ grains; rigors, frequent and feeble heart beats, increase of temperature, increase of secretions, distress, and affections of sight, paralysis of lower limbs, intense headache, no loss of consciousness or sensation, diminution of organic changes. The paralytic action is at first confined to the extremities of the motor nerves; in very large doses this effect extends to the intra-vascular extremities of the vaso-motor nerves. Death occurs from paralysis of respiration. Elimination is rapidly effected by the kidneys. The physiological effects of curare and curarine are identical, but the latter is the more powerful.

Antidotes.—Ligature above the wound by which the poison has been introduced; artificial respiration.

As the greatest uncertainty prevails as to the nature and source of the ourari used, it is quite essential to test that which it is proposed to administer. We have therefore to ascertain (1) the genuineness of the ourari (2) the degree of its activity—for the samples are never alike. It has been employed in tetanus (Vulpian, Follin, Lochner, Neudorfer, and others); epilepsy (Mandl, Benedict, Kunze, Du Cazal, Edlefrei, Bourneville and Bricon); chorea,

strychnine poisoning (Richter, Burow, Corona); convulsive *tic* (Gualla); meningitis (Landerberger). Curarine has only exceptionally been employed, by Busch and Reigel among others; the latter injected in a case of epilepsy, without effect, up to a dose of over $\frac{1}{6}$ of a grain.

"According to Waring, the dose of curarine hypodermically is $\frac{1}{200}$ to $\frac{1}{100}$ of a grain. He says: 'That ourari controls the spasms of *tetanus*, is beyond question, but how far it is really effectual in promoting an actual cure has yet to be shown. The best mode is to inject hypodermically grain $\frac{1}{10}$ in aqueous (filtered) solution of the strength of 1 in 100.'" (Currie).

In injecting subcutaneously, it is best always to select a limb, in order not to lose the resource of ligating in case of accident. Among the local effects observed are: Swelling, pain, local and general rise of temperature, considerable redness, subcutaneous puffing, rosy-white elevation like urticaria (Voisin and Liouville, Du Cazal); painful swellings followed by abscess which persist perhaps several months, probably from the use of unfiltered solutions.

R Ourari, 1 grain,
Hydrochloric acid, 5 minims,
Distilled water to make 15 minims.

Dose 5 drops ($\frac{1}{10}$ gr.) carefully increased.

R Curarine, $\frac{1}{10}$ grain,
Cherry laurel water, 60 minims.

Dose 5 to 10 minims ($\frac{1}{200}$ to $\frac{1}{100}$ grain).

It does not appear to cause pain when administered hypodermically, but may induce slight induration.

℞ Pure emetine, 3 grains.
Sulphuric acid aromatic, 3 minims.
Distilled water, up to 48 minims.
Filter.

Dose, 1 minim ($\frac{1}{10}$ grain).

The human subject can tolerate considerable quantities of it, especially if injected in the subcutaneous cellular tissue; even to $3\frac{1}{2}$ grains introduced several times. It is possible to obtain no poisonous effect from it—not even any inconvenience. The doses indicated as emetic by the hypodermic method by M. d'Ornellas are, on the other hand, very small, viz. from $\frac{1}{16}$ to $\frac{1}{4}$ grain. Foussagrives (*Formulaire Thérapeutique*, 1882) recommend a dose of $1\frac{1}{4}$ grains hypodermically; Dyce Duckworth, $\frac{1}{4}$ grains.

In doses of $\frac{1}{4}$ to $1\frac{1}{2}$ grains, it induces burning taste, salivation, nausea, vomiting, diarrhœa, profuse sweats. During the period of nausea and vomiting, first quickening, then slowing, of the heart-beats and of respiratory movements (Ackerman), with fall of temperature. In frogs there is loss of spinal reflex, irritability and paralysis of the peripheral motor nerves, and of the muscles (Weyland, Pécholier, Harnack).

In poisonous doses.—Extreme muscular debility, collapse and death ($\frac{2}{3}$ grain in a cat and $1\frac{1}{2}$ to $4\frac{1}{2}$ grains in a dog).

In emetic doses.—One-fourteenth to two-sevenths

of a grain (Nothnagel and Rossbach). According to Gubler, it is necessary to give not less than 3 to 4½ grains to produce the same effects as are obtained by 16 to 24 grains of powdered ipecacuanha.

ERGOTUM. ERGOTINUM. ERGOTININUM.
ACIDUM SCLEROTICUM.—EXTRACT
OF ERGOT. ERGOTIN. ERGOTININ.
SCLEROTIC ACID.

[All preparations of ergot when employed hypodermically, cause pain and inflammatory symptoms, unless the injection is made deep into the muscular tissues, when there are no unpleasant local after-effects.

So-called ergotin and ergotinin are delusions, being merely *extracts* of ergot, hence the latter and sclerotic acid are all that demand attention.

℞ Purified ergotin, (ext. ergot), P., D. & Co.'s. Dose, 2 to 6 minims in water; represents 10 to 30 grains of ergot.

℞ *Normal liquid* ergot, P., D. & Co.'s (assayed), 5 to 30 minims, at one dose.

℞ *Acid sclerotic*, P., D. & Co., 18 grains.

Chloral hydrate, 3 grains.

Camphor water, 1 fluidounce.

Dose, 5 to 30 minims *plus*.—ED.]

ETHER.

Sixteen to thirty-two minims may be used hypodermically; is sometimes employed as a diffusive stimulant. It may be diluted with water, 1 to 9.

Spirits of sulphuric ether, (Hoffmann's Anodyne), and acetic ether, have been employed as analeptics. It will be well to remember that ether dissolves the cement by which the joints of the average syringe are secured.

EUCALYPTOL.

Eucalyptol is soluble in alcohol, sparingly so in water.* It is an anti-fermentescient and anti-putrescent, and produces, in doses of from 32 to 64 grains, cephalalgia, physical and mental prostration, loss of reflexes, failure of respiration, fall of blood pressure and temperature. *In poisonous doses*, death occurs from arrest of respiration. *Therapeutics.* --Eucalyptus and its derivatives have been often used successfully, both internally and externally, in a great variety of diseases, but eucalyptol has only been administered subcutaneously by Dr. Sloan, who states that he has saved thereby the lives of patients suffering from severe puerperal fever (Kobner).

FERRUM.—IRON.

Dialyzed iron, diluted and undiluted, has been used by Da Costa in doses of 15 minims in cases of anæmia. M. Luton has injected it in doses of 3, 5, and 10 minims: "Deprived of its styptic and astringent properties, it is nearly as well borne by the cellular tissue as perfectly neutral solutions of morphine,

* See Vaselin, page 122.

is rapidly absorbed, and locally gives rise to nothing more than a trifling nodosity, which, however, persists for a considerable time. The immediate effects are similar to those of an energetic diffusible stimulant; a sense of warmth more agreeable than painful is diffused throughout the body, the arteries pulsate strongly, the countenance is flushed, there is increased mental activity, rapidity of thought, and desire for exercise. These effects last some time, proportionately to the dose injected, and are never followed by prostration, but by a certain degree of anorexia." M. Luton has also observed a re-awakening of the generative desire.

The *pyrophosphate of the ammonio-citrate* has been employed by Hugenin, Neuss, Lippert, and others, but it is by no means certain that it is not apt to induce abscess.

℞ Pyrophosph. of ammon.-cit. iron, 3 grains.

Distilled water, up to 16 minims.

Sixteen minims equal approximately $\frac{1}{3}$ grain of iron.

This solution should be made as required, and kept in a cool place. Iron is detected in the urine thirty minutes after injection.

GELSEMIUM. YELLOW JESSAMINE.

Gelsemium paralyzes the motor centres of the brain and the respiratory centres (Berger, Moritz, Ott); it first increases then diminishes reflex excitability of the spinal cord and the cardiac contractions.

These symptoms are complicated with weight of head, vertigo, diplopia, impeded movements of the tongue, tremor of the hands, numbness of the fingers, nausea, vomiting, dyspnœa, feeling of general cold (Berger). The medicinal doses range from $\frac{1}{100}$ to $\frac{1}{40}$ grain of GELSEMINE, and from 1 to 5 drops of the *normal liquid*.

Hydrochlorate of gelsemine, P. D. & Co., 1 grain.
Distilled water, up to 210 minims.

From 2 to 5 minims, equivalent to $\frac{1}{140}$ to $\frac{1}{70}$ of a grain of salt; it produces insensibility at the point of injection.

[The *normal liquid* is every way preferable to the alkaloid, and may be given diluted with glycerin, alcohol, or glycerin and camphor water.—ED.]

It has been successfully used in the treatment of neuralgia by Drs. Jurasz, Wickham Legg, Thomson, Clarke, G. A. Stockwell, and others. The hydrochlorate is ten times as active as the *gelsemin*, or concentration, and is a permanent salt.

HYDRARGYRUM—PREPARATIONS OF MERCURY.

Bi-chloride of mercury is soluble in 6 parts of cold water, and freely soluble in alcohol, ether, and glycerin. *Cyanide* of mercury is soluble in water (5 per cent.), and dilute alcohol; slightly soluble in strong alcohol, and in glycerin. *Bin-iodide* of mercury is insoluble in water and in glycerin, soluble in al-

cohol and in ether. In *small doses*, according to some authors, mercury is stimulating and restorative as well as antiseptic. In medicinal doses it has no appreciable effect on the formation of urea, but it is generally admitted that its *prolonged* use is attended, with diminution of globulin.

Physiologically it produces stomatitis more or less speedily according to idiosyncrasy or local predisposition, salivation, intestinal catarrh, diarrhœa; often notable slowing of the circulation and fall of temperature. Mercurial fever is generally due to local irritations produced by these preparations (Barrallier). When pressed to procure chronic mercurialism, it is a brain poison, one of the most constant phenomena being a condition of timidity and embarrassment, falling out of the teeth, shrinking of the gums, chronic inflammations of the mouth and pharynx, induration of the salivary glands and of the cervical ganglia, gastric catarrh, pains in the limbs, nervous excitability, insomnia, vertigo, tremors, impairment of memory, frequently emaciation, dental neuralgia, headache, dyspnœa, tinglings, anæsthesia and analgesia.

R Corrosive sublimate, 1 grain.
Distilled water, up to 168 minims,

Dr. Lewis injected 12 minims of this solution, to which he sometimes added morphia and glycerin. Or the formula may be made:

is greater contrariety of opinion than that of the utility of subcutaneous injections of mercury for syphilis. Dr. Buzzard concludes it is safe with ordinary care as regards the formation of abscesses, and that the necessary mercury is absorbed into the system. He believes it, however, to be a painful mode of treatment, even when the injection is made in the back, and the solution employed quite clear. He is disposed to think that it possesses no advantages over mercurial inunctions in cases of gummatous syphilis, or when there is evidence of internal disease, except perhaps the very problematical one of substituting a clean but painful process for one which is as dirty as it is free from personal suffering. In this opinion Messrs. Hill and Cooper coincide.

Bäumler, while admitting that subcutaneous injections made with due precautions, beyond the pain which they occasion, need give rise to no local troubles except in unhealthy persons, thinks that the method is not so rapid in its effects as appears at first sight, the rapid disappearance of the external symptoms of syphilis being by no means equivalent to a cure of the disease itself. He also quotes the observation of Zeissl, to the effect that there is more apt to be recurrence of syphilitic manifestations after hypodermic than any other form of medication. Later observers, among others Villar, Florez, and Schultz, speak highly in praise of anti-syphilitic hypodermic injections. The two former authorities recommend

injections of mercuric peptone and of cyanide of mercury, and state that the latter salt is especially useful in cases of ocular, cerebral, and medullary syphilis. They claim that hypodermic medication possesses the advantages over treatment by internal administration or by inunction, of being more exact in regard to dosage, in not giving rise to gastro-intestinal troubles, in shortening the duration of treatment, and in rarely causing pyalism. Schultz uses a combination of mercuric *chloride* with *urea*.

Scarenzio, Ambrosoli, and Sigmund, used hypodermic injections of *calomel*, by means of a finely levigated powder held in suspension in water and glycerin, in various syphilitic affections. Maximum dose, $\frac{3}{4}$ grain.

Calomel, 3 grains.

Glycerin, up to 16 minims.

M. G. Smirnoff treated syphilis successfully by hypodermic injections of calomel. He employed solutions of one-half the strength of those employed by Scarenzio and others, with the object of avoiding abscess.

"M. Boni was much pleased with the results obtained by injections of calomel in croup; $\frac{3}{4}$ grain in the left arm of a child" (Currie).

HYDRASTININA.—HYDRASTININE.

[Hydrastinine in pure condition is perfectly white, and melts in the test tube at 116° to 117° C.; is soluble in alcohol, ether, and chloroform, with extreme

facility; more difficultly soluble in warm water, and with most of the acids readily forms soluble salts. The *hydrochlorate* is readily soluble in water; the solution exhibits slight fluorescence, possessing, with the free base, an intensely bitter taste. In both warm and cold-blooded animals, it paralyzes by means of its action on the motor sphere of the spinal cord; a tetanic stage is not produced. It increases the contractile power of the heart muscle. *It is not a cardiac poison.* It has no local effect on the muscular tract. Vascular contractions are produced by its action on the vessels themselves, resulting in increased blood-pressure, which, in the beginning, is periodic, quite pronounced, persistent, and not interrupted by any symptoms of relaxation. Concurrently with the increase of blood-pressure, is observed an irritation of the vagi, which produces a slowing of the pulse. At death, the lessening of blood-pressure is of a secondary nature not dependent upon a relaxation of the blood-vessels, and removable by means of artificial respiration. Death ensues in consequence of paralysis of the respiratory centres.

The above, showing the absence of irritation of the spinal nerves, the favorable action upon the heart, and persistent contractility of the vascular system, is convincing that hydrastinine is better suited for therapeutical employment than hydrastine—the active principle of *Hydrastis Canadensis*. On the other hand, the absence of an action on the mus-

cular portion of the tissues, is a powerful argument for the subcutaneous injection of this remedy.

Dr. Falk, after an injection in his own person of 1 gramme (16 drops) of a five per cent. solution, subcutaneously in the fore-arm, was convinced that it is not any more painful than cocaine. Directly after its use, in the neighborhood of the puncture there is a feeling of cold and insensibility, otherwise nothing either subjective or objective; never any abscesses, though it is not always possible to avoid infiltrations that cause temporary swelling. These local effects disappear speedily, however, being disconnected with any sensation of pain.

Dr. Falk summarizes his experiences with the product as follows:

In cases of metrorrhagia, wherein hydrastinine was employed, in which anatomical lesions were absent, the remedy established its usefulness decidedly. In congestive dysmenorrhœa, menstruation progressed during its use without the occurrence of congestive troubles; the hæmorrhage was much decreased, and the period shortened. Equally good results were obtained in the metrorrhagia of virgin uteri. In all cases in which menstruation occurred every fourteen days and with a duration of from eight to twelve days, hydrastinine shortened the period to five days, with an interval of from five to five and a half weeks, with a notable increase of strength.

In cases of pyosalpinx, hydrastinine was very

serviceable, relieving the congestions of the genitals and decreasing the amount of hæmorrhage. In cases of metrorrhagia due to endometritis, and, in those complicated with chronic metritis, there was decided improvement. In all cases menstruation had been previously very irregular; in several, intercurrent hæmorrhages, in others menstruation, occurred at intervals ranging from fourteen days to three weeks, lasting for a long time, naturally weakening the patient by loss of blood and other concurrent causes. These hæmorrhages had been treated by repeated abrasion of the uterus, and by ergotin, but without result. In all, the intercurrent hæmorrhages desisted, menstruation became regular, and lasted but a short time, with a notable improvement of attendant symptoms. Women who usually were too weak to perform any work during the menstrual period, declared that after the injections they had gained sufficient strength to indulge in their usual avocations, and did not require the same amount of rest as formerly; and the improvement was permanent.

Excellent results were obtained in cases of myoma of different characters and sizes, in all of which there was more or less hæmorrhage. After a few injections, the hæmorrhage, which usually lasted from four to eight weeks, and had proven uncontrollable by ergotin, became only a slight loss of blood; and there was a consequent lessening of debilitating influences.

Out of twenty-six patients of the foregoing classes treated with hydrastinine, only four were not either improved or cured. The successful results can, therefore, be summed up as giving 84.6 per cent. of the cases.

A critical summary of cases, leads to the inference that those in which menorrhagia preponderate (also those of congestive dysmenorrhœa), and mild attacks of endometritis, define the boundaries of this treatment. It is not unlikely also that hydrastinine can exert a contractile effect on the vessels, and by that means influence the character and quantity of hæmorrhage; and it is also possible that it is an excitant or stimulant to the muscular structure of the uterus. In evidence of this position, may be adduced the fact that pains resembling those of labor, and slight hæmorrhages, ensue after its injection, principally in cases of endometritis.

Dr. Falk frequently began its use during hæmorrhage, and found in cases of endometritis even after the first injection, a cessation of the flux; also in cases of myoma, after several injections. The most satisfactory results were, however, obtained in cases of congestive dysmenorrhœa, or too profuse menstrual hæmorrhage dependent upon structural changes of the uterus, when the treatment was adopted from six to eight days before the expected occurrence of the menstrual flow. In cases of irregularity, he injected every other day, or perhaps two to three times a

even if accompanied by depression; in ordinary epilepsy; and in delirium tremens. Currie says henbane lessens the delusions and cuts short the attacks. He adds "that in all cases of mental excitement, including acute mania, hysterical mania, and puerperal mania, it is the only reliable hypnotic that we have."

HYOSCINE is a somewhat glutinous material, and yields crystalline salts, the principal of these being the *hydriodate*, the *hydrobromate*, and the *hydrochlorate*. Experiments have been made with the two former, the hydrobromate being mostly used [and every way the most preferable as it is had in permanent colorless crystals.—ED.]

Hyoscine acts as a spinal, and centric respiratory depressant; it affects the circulation slightly, and does not paralyze the pneumogastrics. In poisonous doses, although it paralyzes the vaso-motor system, it has very slight depressant influence on the heart. In the healthy human being, the effects of a medicinal dose are: Ten minutes after subcutaneous injection the symptoms resembled those of slight alcoholic intoxication, viz., sense of lightness and giddiness; these were succeeded by dryness of the mouth and throat, and restlessness, in spite of giddy sensations. Then followed drowsiness and profound sleep, on awakening from which there was no recollection. In one gentleman who accidentally took a small dose of a strong solution, there were symptoms of weight of the eyelids, constrictive headache (chiefly frontal), sense of suffocation without dyspnœa, great uneasiness and

unrest, with intervals of unconscious sleep. On recovery, the after-effects were limited to slight headache and feeling of lassitude. It is of little use to relieve pain, but of great value in spasms, and is contraindicated in patients with a tendency to pulmonary œdema. Also is of great service in the treatment of delirious excitement and insomnia, in insanity, and other diseases; has been found useful in controlling the delirium of pneumonia, cardiac and renal disease, epileptiform convulsions and chorea. Prof. Edlessen administered it in six cases of whooping cough, with advantage in three; in six cases of asthma, always with advantage; and in two cases of severe enteralgia, with relief of pain. It has been extensively used by alienists with unmistakable benefit in cases of acute dementia, mania, melancholia, hystero-epilepsy, and other mental diseases. It is, however, "only a temporary remedy to quiet delirium and control excitement, and does not cure the various diseases in the course of which delirium arises" (Currie).

It is best administered hypodermically, the doses ranging from $\frac{1}{800}$ to $\frac{1}{100}$ of a grain, and in some cases to as much as $\frac{1}{75}$ or $\frac{1}{50}$ of a grain. Dr. Mitchell Bruce has given as much as $\frac{1}{3}$ of a grain of the hydriodate without injurious effects, but as a general rule this dose is unsafe, and even with $\frac{1}{100}$ of a grain Dr. Nestor Tirard observed, in a case of chronic Bright's disease, angry and combative delirium.

℞ Hydrobromate of hyoscine, 1 grain.
Camphor water, 200 minims.

Dose, 1 minim = $\frac{1}{800}$ grain.

ICHTHYOL.—SULPHO-ICHTHYOLIC ACID.

A brownish yellow oil soluble in water, and recommended by Unna in the treatment of skin diseases; he also recommends it internally for acute and chronic rheumatism. Although chiefly used as an external application, it has been employed also subdermically, in 16 minim doses of a 5 to 10 per cent. aqueous solution. [After all, ichthyol is simply a very soluble preparation of *sulphur*; being a powerful antiphlogistic, rapidly producing anæmia of all the tissues to which it is applied, it must not be permitted to touch abraded surfaces. Unna says it is indicated internally in doses of from 30 to 60 grains.—Ed.]

ODOFORMUM.—TER-IODIDE OF FORMYL.

Insoluble in water, soluble in alcohol (1 in 50), in ether (1 in 6), in fats, volatile and fixed oils, vaselin, chloroform, and carbon sulphide.

The ethereal solution of iodoform has been used with success in severe cases of arthritis, by Messrs. Miculicz, Neumann, Müller, etc. Elimination is slow, yet its presence is speedily detected in the urine. 1½ drachms have been injected in thirteen days. It has been successfully employed in syphilis, but chiefly internally and externally in goitre, scrofula, rickets, phthisis, anal fissure, prostatic enlargements, neuralgia (Moretin), uterine and mammary carcinoma (Demarquay), serous collections, chronic arthritis, glandular enlargements (Gubler), cancerous and vener-

eal ulcers (D'Amico), to stimulate sluggish wounds, in varicose and phagedenic ulcers, rupia and ecthyma, inter-digital mucous gummata, suppurating buboes, moist gangrene, gunshot wounds (Féréol), intermittent fevers (Knotts), toothache due to caries (Lailier, Marrotte.)

℞ Iodoform, 6 grains,

Glycerin, up to 24 minims.

From 4 to 10 minims of the solution.

(Thomann.)

℞ Iodoform, 1 grain,

Almond or castor oil, up to 16 minims.

The effect subdermically is, local and general anæsthesia, antisepsis, vomiting, nausea, epigastric pains, diarrhœa (Maillard, dose 7 grains). [The fact is, it is of little value subdermically.—ED.]

JABORANDUM.—JABORANDI.

[Hypodermic injections of PILOCARPINE salts, or even of the *normal liquid* jaborandi properly diluted, so far as the writer can discover, has never been attended with any accident or untoward result not due to the physiological activity of the drug exclusively.—ED.] The physiological phenomena include congestion of the skin, principally of the face; increase of secretions, especially of the salivary and sudorific glands; acceleration of cardiac pulsations; initial increase of temperature, sometimes persistent if diaphoresis is not established, with consecutive fall in accordance with the greater or less abundance of perspiration. In medic-

inal doses the salts of pilocarpine are less apt to produce nausea, vomiting, diarrhoea, and that sense of extreme lassitude, which frequently result from the use of jaborandi; and it may be observed that in doses of less than $\frac{1}{4}$ grain the unpleasant effects resulting from administration of the salts, have for the most part been observed in patients suffering from cardiac affections.

R *Muriate* (or *nitrate*) of pilocarpine, 2 grains.

Distilled water, up to 160 minims.

Dose, 5 to 10 minims *plus* ($\frac{1}{4}$ to $\frac{1}{2}$ grain).

In is usually best to begin with a small dose and increase cautiously; when it exceeds $\frac{1}{4}$ grain, it is advisable to administer in two or more portions—one-half in the morning, the other in the evening. [I have never given hypodermically more than $\frac{1}{2}$ grain, though Bourneville and Bricon admit employing $\frac{3}{4}$; the former dose sometimes secures profound exhaustion. One-eighth grain introduced in the arm during the cold stage of intermittents, invariably, in my hands, aborted both chill and fever, sweating at once supervening; and I have found both pilocarpine and jaborandi invaluable in treating obstinate (chronic) cases of ague.—ED.]

Injections have been specially used in the treatment of various forms of dropsy, in pleurisy, bronchitis, pneumonia, acute articular rheumatism, epilepsy, eclampsia, mercurial poisoning (Prokop, Rokitansky), unilateral sweats (André), foetid perspira-

tion of the feet (Armaingaud), rapidly progressive posterior sclerotico-choroiditis with muscæ volitantes and threatened detachment of the retina (Dehenne, Deniau), detachment of the retina (Dianoux and others), some forms of polyuria (Roques, Huchard and Ducroux), syphilis (Lewin), diphtheria (Lehwess and Guttmann); [hypopion, Henry Williams, C. B. Stockwell—ED.]; also to assist uterine contractions; in uræmic coma, asthma, puerperal convulsions, cholera; and to relieve the pains of tabes dorsalis. "An interesting case of cure of alopecia from pilocarpine injections is recorded in the *Lancet*, vol. i., 1882. Brünauer has used it with success in a case of tetanus of rheumatic origin; M. Chéron has injected $\frac{3}{4}$ of a grain daily as a galactagogue, with complete success in eight out of nine cases; Dr. Purjeck cured a case of atropine poisoning, and Dr. Roth, a case of stramonium poisoning, by hypodermic injections of pilocarpine." (Currie.)

There is an unmistakable antagonism between pilocarpine and belladonna or atropine.

MOSCHI TINCTURA.—TINCTURE OF MUSK.

M. Rosenthal praises musk highly as a diffusible stimulant in the agid stage of cholera, and in adynamic forms of typhoid fever. M. Rhode used an aqueous solution (1 in 40), of which he injected from 16 to 48 minims *per diem*, according to the gravity of the case. In most of the cases a rise of pulse-rate and of temperature resulted.

Locally, he observed on one occasion a slight slough of the skin in a case of cholera, and in another instance there was induration without abscess. Musk and its tincture have lately been used by various writers, among others by Vogelsang, in the treatment of infantile convulsions. Tincture of musk, made in the proportion of one part of pure pod musk and twenty-five parts each of alcohol and water, is injected in doses of from 5 to 30 minims.

MUSCARINA.—MUSCARINE.

Has been employed hypodermically by Ringer and Morehead, who gave in doses of $\frac{1}{3}$ to $\frac{3}{4}$ grain. M. Prevost has suggested that it might be possible to antagonize the toxic symptoms due to atropine by hypodermic injections of muscarine, physostigmine, or pilocarpine, in moderate doses; he proved a natural antagonism between muscarine and atropine by experiment. It has been used with success in the night-sweats of phthisis, and to arrest the secretion of milk. [Muscarine *sulphate* is very hygroscopic; dose hypodermically $\frac{1}{100}$ to $\frac{1}{8}$ grain.—ED.]

NICOTINA.—NICOTINE.

Nicotine was employed hypodermically by Erlenmeyer in a case of tetanus, without local trouble and with good results, in doses of $\frac{1}{10}$ of a grain. [Is an excellent antagonist to strychnia.—ED.]

R Nicotine, 1 grain.

Chloroform water, up to 280 minims.

Dose, four to eight minims = $\frac{1}{10}$ of a grain.

NITRO-GLYCERINUM.—GLONOID. TRINITRIN.

An inodorous, yellow oil of mawkish taste, but slightly soluble in water, though readily so in alcohol or ether. It is a powerful poison, and in moderate doses (1 to 6 drops of a 1 per cent. solution) produces a feeling of intercranial fullness, more or less severe headache, confusion or sluggishness of thought, tinnitus aurium, vertigo, amblyopia, quickening of pulse rate with occasional diastole, diminution of arterial pressure, redness and flushing of face, sweats, nausea, vomiting, and occasional exaggeration of the respiratory movements. In *poisonous* doses (excess of six drops) clonic and tonic convulsions succeed, followed by death from asphyxia. The effects closely resemble those of amyl nitrite, but are more persistent.

Glonoïn has been administered internally in neuralgia, epilepsy, and hysteria, by Thorowgood, James, Laurence, Baker Edwards, Brady, and others. In consequence of the experiments of Fuller, Harley and Vulpian, who disputed the results obtained by the above mentioned authors, glonoïn fell into disrepute for several years, but after the publication of M. Bruel's essay, was again used by Murrell, Mayo-Robson, Craig, Farquharson, Stitts, McCall Anderson, Green, Hammond, Stewart, Korcinski, Huchard, Trusevich, and Bramwell, in angina pectoris, acute and chronic nephritis, epilepsy, eclampsia, asthma,

palpitation, affections of the heart and aorta, syncope, migraine, neuralgia, vertigo, Menière's disease, and epileptiform *tic*. "In addition it has been employed with marked benefit in seasickness, the vascular tension which occurs in the aged, myxœdema, and puerperal convulsions" (Currie).

In regard to some of these ailments, the results obtained are contradictory. Glonoin appears to be mostly of use in angina pectoris, and in those diseases in which cerebral anæmia is a symptom. Quite recently M. Rossbach has recommended its use subdermically in the treatment of interstitial nephritis.

R Nitro-glycerin solution, 1 per cent., 5 drachms.

Rectified spirit, 2 drachms.

Distilled water, up to 1½ ounces.

Dose, 1 to 4 minims. One minim contains $\frac{1}{16}$ of a grain.

NUX VOMICA.

Nux vomica increases the tactile sensibility, causing tinglings, salivation, anxiety, vertigo, rapid breathing, and stiffness of nuchal muscle; tetanic spasms in rapid succession, merging into one another, with opisthotonos and forcible extension of the lower limbs and of the trunk; trismus from the action upon the bulb and spinal marrow; dilatation of pupils, elevation of blood pressure, abolition of energy of the motor nerves, slowing of cardiac pulsations; death by asphyxia, arrest of heart, and unascertained modification of the nerve centres.

STRYCHNINE is very sparingly soluble in water, alcohol and ether, but its salts are all soluble, and more toxic than the alkaloid; the *muriate* and *nitrate* are more active than the *sulphate*. Strychnine is also possessed of decided antiputrescent and antifermentative properties, and is not generally admitted to possess cumulative action. The salts administered hypodermically do not cause any local trouble save that sometimes erection of the hair follicles (goose skin) is observed at and near the point of puncture.

According to Husemann, the minimum lethal dose of strychnia is $\frac{1}{2}$ grain. It should at first be administered in very minute doses ($\frac{1}{125}$ to $\frac{1}{60}$), cautiously increased until the appearance of toxic phenomena (stiffness of muscles of neck, spasms, etc.), when the administration must be suspended. *Antagonisms*: Calabar bean (Watson, and Keyworth), chloral (Liebrich, Rajewsky), chloroform inhalations (?). A pseudo-antagonism exists between strychnia and curarine. [Dr. Currie remarks: "Urethan is alleged to be nearly as efficient and much less dangerous an antagonist than chloral. Dr. Bignon concludes, from experiments made upon dogs, that cocaine is a physiological antagonist to strychnine; up to a certain point after the ingestion of poisonous doses, the animal could always be saved by hypodermic injections of cocaine, but when too large doses were administered it always perished from the correspondingly large doses of cocaine required." —ED.]

℞ Nitrate of strychnine, 1 grain.

Distilled water to make 300 minims.

Dose, 5 minims ($\frac{1}{60}$ grain).

℞ Sulphate of strychnine, 1 grain.

Dilute sulphuric acid, 3 minims.

Distilled water to make 300 minims.

Dose, 5 minims ($\frac{1}{60}$ grain).

Nux vomica and strychnine have been recommended in the treatment of delirium tremens; and Dolbeau advises $\frac{1}{10}$ of strychnine sulphate injected around the anus *in divided doses*, every two days, in the rectal prolapse of children. Luton advises:

℞ Sulphate of strychnia, 1 grain,

Cherry laurel water, 105 minims,

Distilled water, up to 210 minims.

Fifteen minims = $\frac{1}{4}$ of a grain: an injection every half hour until four have been given, after that every hour; he says the dose should reach $\frac{3}{4}$ to $\frac{1}{2}$ of a grain. Parke, Davis & Co.'s hypodermic tablets contain either $\frac{1}{100}$ or $\frac{1}{50}$ of a grain as desired.

Tincture of nux vomica has been employed hypodermically in threatened asphyxia from catarrh, in the death agony, and in the last stages of adynamia, in doses as high as 80 minims (Luton), with no local effects save slight swelling, heat, and moderate pain, all of which soon disappear. [*Normal liquid* is preferable to the tincture however, and oftentimes to the alkaloidal salts, for hypodermic use, and may be employed in doses of from 1 to 5 minims.

BRUCINE, another alkaloid of *nux vomica*, has in general the same physiological action as strychnine, with local anæsthetic effect superadded. It might be employed with advantage in combination with many preparations that cause intense pain when administered hypodermically, being preferable for this purpose to cocaine. It is soluble with the aid of heat in dilute alcohol, and may be given subcutaneously in doses of from $\frac{1}{80}$ to $\frac{1}{20}$ of a grain.—ED.].

All these preparations have been employed in: Paralysis of the vocal cords, of the crico-thyroid muscles (Waldenburg, Neudorfer), saturnine palsy (Lorent), hemiplegia, paraplegia, paralysis from compression of radial nerve, Bell's paralysis, hemeralopia, amaurosis, writer's cramp, urinary incontinence, intestinal atony, lightening pains of tabes, [brucine is preferable.—Ed.], sciatica, pulmonary emphysema, cholera, chloral poisoning, diphtheritic paralysis, infantile paralysis, progressive muscular atrophy, and locomotor ataxia. Maestro Perez suggests injections of 4 to 5 minims of a two-per-cent. solution in the second stage of cholera.

[“Atropine has been employed to antagonize the poisonous effect of strychnine; two cases were successfully treated by Dr. Roberts, in one of which the strychnine was administered in whiskey. Gubler says that possibly atropine is antagonistic, but Bartholow strongly combats this view, asserting the atropine only intensifies the effect of the poison, and hastens

the tetanic fixat
... confirm
... sees that am
... idea that is pr
... experiments
... H. A. Hare
... that its action
... employed to
... can be
... chloral, all hav
... evidence is no w
... It would be
... as its action
... permanent.—
... hypoc
... value in ca
... of Dr. T. I
... this drug stim
... and the former
... volume on the
... similar to its ac
... centres. Dr.
... this respect
... monia, and cl
... stimulants.
... succeeds w
... mends the
... following
... of the

stages (prodromata) of acute infectious diseases; when administered in the initial stage of infectious fevers, it overcomes the symptoms of general malaise, muscular lassitude, etc., and the patient experiences relief even after a single injection. Dr. Russell has found the drug of special use in the treatment of typhoid fever, which disease, he asserts, it aborts in the early stages, or simplifies when administered after the fever is fully developed; and he attributes these effects to the power which the remedy possesses of 'sterilizing, as it were, the soil on which the pathogenetic micro-organisms of typhoid fever vegetate.'" (Currie.)

OLEANDRINA.—OLEANDRINE.

This alkaloid, extracted from the leaves of *Nerium o'leander*, is one of the group of cardiac poisons, and has been administered subcutaneously by Erlenmeyer without success in epilepsy.

℞ Oleandrine, 10 grains.

Alcohol, q. s.

Distilled water to make 120 minims.

Dose, 12 minims.

OLEUM AMYGDALÆ. OLEUM OLIVÆ—AL-
MOND OIL. OLIVE OIL.

According to M. Binz, these oils have been used subcutaneously as nutrients, but without success on account of difficult absorption by the cellular tissue. [Shoemaker, of Philadelphia, claims to have

used it in a number of cases with encouraging results, but his statements are to be accepted *cum grano*.—ED.]

Krueg has injected four to eight drachms of olive oil per diem in the case of a lunatic who refused food. The injection fluid being expelled slowly, caused no pain, and left only a trifling redness.

OPIUM.

[Opium has a much more decidedly stimulating effect than morphine on the cardio-vascular apparatus; it raises the pulse, at the same time increasing the amplitude of the pulse wave, and it stimulates also, in the most marked way, the calorific function. The *normal liquid* of Messrs. Parke, Davis & Co., is an excellent preparation for hypodermic use:—ED.]

MORPHIA alkaloid is almost entirely insoluble in water, sparingly soluble in alcohol, ether, chloroform, and glycerin, and insoluble in essential oils.

[The *hydrochlorate* is soluble in 16 to 25 parts of water, 50 parts of alcohol, and to the extent of 20 per cent. in glycerin. The commercial *sulphate* is an undesirable salt for hypodermic use, and is less soluble than any other form, the *meconate* excepted. The *acetate* ranks next to the hydrochlorate, and contains 80 per cent. of morphia alkaloid to only 76 found in the sulphate.—ED.]

In medicinal doses morphine causes somnolence, muscular enfeeblement, sensorial troubles, contraction

... mental torpor; diminution of sensibility ... muscular contractability co-existing with in- ... reflex excitability; dryness of mouth and ... thirst, impairment of appetite and of gas- ... constipation, nausea, vomiting; more ... perspiration; itchings, excitement ... awakening. In larger doses there are ... in thought and movement, succeeded by ... sleep; the pulse is either slow or of un- ... frequency; respiration becomes irregular, and ... falls in number to eight or ten per minute; ... membranes become cyanosed; the counte- ... pallid, and covered with violet colored ... the temperature falls; and death finally ... frequently amidst clonic and tonic ... *Morphinism* produces dryness of the mouth, ... nausea, vomiting, generally constipation, occa- ... dyspnoea and palpitation [a combination rare ... Ed.], albuminuria, loss of power, amenor- ... insomnia with hallucinations, varying disposition, ... aesthesia, neuralgia, tremor of hands, bewildered ... stupid expression; various forms of eruptions, ... appetite, tendency to suppuration, emaciation; ... phenomena are more prominent after the sud- ... abandonment of the morphine habit.

Man is, of all animals, [the dog excepted,—Ed.] ... most sensitive to the action of morphine, hence it ... necessary to be on guard against the error of ... adding the effects likely to arise. Besides, in man

the effects vary with age, constitution, etc. Children, for instance, are extremely susceptible to the influence of morphine. The lethal dose in individuals who are not habituated to its use, varies within very wide limits; hence with such it is wise to begin with a minute dose ($\frac{1}{16}$ grain for example), and increase cautiously. Morphine is *eliminated* with considerable rapidity by the kidneys. The *local effects* are *nil* when the injection is skillfully made.

Solutions of morphine hydrochlorate in water alone, become turbid very quickly, and form *apomorphine* with irritant properties. To obviate this difficulty, waters distilled from eucalyptus globulus (Gubler), peppermint, and cinnamon, have been proposed, (Delioux and Savignac); also glycerin, alcohol, alcohol and water (Adrian and C. Paul), distilled elm-water (Patrouillard), weak chloral solution (Vidal), a solution of salicylic acid, 1 in 2,000 (Limousin, Keyes, etc.), carbolic acid; cherry laurel water also has been employed, but it rapidly oxydizes by exposure to the air, when the solution causes pain.

℞ Morphine hydrochlorate, 12 grains.
Chloral, 15 grains.
Camphor water, up to 480 minims.

—ED.]

℞ Morphine hydrochlorate, 1 grain.
Chloroform water, 40 minims.

Five minims of either of these solutions represents $\frac{1}{8}$ grain.

—ED.]

~~... of morphine, 12 grains.~~

~~... 2 minims.~~

~~... water to make 480 minims.~~

~~... grain. Ed.]~~

~~... of morphine, 12 grains.~~

~~... 1 minim.~~

~~... hydrate, 2 grains.~~

~~... glycerin, 1 drachm.~~

~~... water, q. s. to make 480 minims.~~

~~... morphine after mixing with the acetic acid,~~

~~... of water, then add chloral hydrate, and last~~

~~... mixed with balance of water; filter—5 minims~~

~~... —Ed.]~~

~~... morphine has become so extensive~~

~~... that it is administered now-a-days in all~~

~~... its rightly and wrongly. It is impos-~~

~~... too strongly against this abuse, as~~

~~... and its salts should never be prescribed~~

~~... defined cases. Morphine injections~~

~~... been used in the treatment of cholera.~~

~~... Coles, recommended morphine before the~~

~~... collapse, combined with atropine in the~~

~~... stage, and atropine alone in the stage of~~

~~... See Belladonna.)~~

~~... a pity hypodermic tablets of morphine mur-~~

~~... acetate are not in the market.—Ed.]~~

~~... VERINE has been employed hypodermically~~

~~... Leidesdorf and Schüle, in cases of insanity,~~

~~... of *hydrochlorate* or *phosphate* (six to twelve~~

~~... of a 1 in 12 solution); the phosphate produced~~

~~... and sloughing.~~

APOMORPHINE HYDROCHLORATE.—This differs from morphine in having one less equivalent of water; the hydrochlorate alone is employed.

℞ Hydrochlorate of apomorphine, 2 grains.

Camphor water, 100 minims.

Dose, 2 to 8 minims (Currie).

A few drops of muriatic acid may be added when concentrated solutions are made, to clear them up. It dissolves with difficulty in cold water, but rapidly in warm; the solution after the lapse of a few moments assumes a green tint, and is apt to be unstable, hence it is best made fresh as required. [Messrs. Parke, Davis & Co.'s hypodermic tablets are invaluable for this purpose.

Hydrochlorate of apomorphine should only be used with the utmost caution in consequence of the peculiar tendency to syncope and collapse which it induces in some subjects. The majority of works on therapeutics, and the formulæ of many authors, give as the emetic dose $\frac{1}{4}$ up to $\frac{1}{2}$ of a grain, but the latter is decidedly dangerous. Indeed no emetic has given rise, (regard being had to the frequency of its use,) to so many cases of collapse and syncope. In one case $\frac{1}{8}$ grain caused great prostration (Currie), and $\frac{1}{16}$ must be deemed sufficient for an adult if of good quality.—ED.]

Its action is directly on the nerve centres, exciting vomiting, and not, as is generally imagined, reflex (David). Feeling of weight at the epigastrium, slight

headache, acceleration of pulse and respiration, salivation, perspiration and nausea, precede emesis. Though chiefly prescribed as an emetic, diaphoretic, and expectorant, it has also been employed in gastric diseases and obstruction, various forms of poisoning;* in threatened asphyxia, croup, pneumonia, tonsillitis, broncho-pneumonia; layrngismus stridulus, foreign bodies in œsophagus, mania, hystero-epilepsy, angina tonsillaris, diphtheria, epilepsy. Hypodermic injections procure no pain worthy of mention, and no local disturbance except possibly small subcutaneous indu-rations which rapidly disappear.

CODEINE.—The *hydrochlorate* of codeine, which is usually employed hypodermically, is soluble in 80 parts of water, in alcohol, and in ether.

℞ Hydrochlorate of codeine, 2 grains,
Distilled water, 160 minims.

Ten minims = $\frac{1}{4}$ grain.

Phosphate of codeine is soluble in 4 parts of water and contains 70 per cent. of codeine; it should be administered in double the dose of morphine. "Codeine has been used in gastrodynia, cancer, and

* As morphine delays the development of the action of apomorphine, and as chloroform and chloral injected intravenously delay the action until awakening occurs, it would be useless—nay more, *dangerous*—to administer hydrochlorate of apomorphine in poisoning by morphine, chloral, and chloroform, owing to the time which would be lost, if the poisoning were sufficient to produce anæsthesia and muscular relaxation (David).

insomnia. It is very useful in some forms of cough, and particularly so in diabetes. In one case of obstinate bronchial asthma, I have found it more effectual than any other drug in quieting the paroxysms and inducing sound refreshing sleep." (Currie). Erlenmeyer employed it in neuralgia without apparent benefit; Reissner in mental diseases in doses of $\frac{3}{4}$ to $1\frac{3}{4}$ grains. Very exceptionally there are inflammatory troubles at the point of injection. [Recently has been widely heralded as a specific in yellow fever; and certainly its physiological characteristics favor the claims made.—ED.]

APOCODEINE and APONARCEINE are little employed.

℞ Hydrochlorate of apocodeine, 3 grains.

Camphor water, up to 224 grains.

Dose, 15 to 25 minims.

The effect is produced in a few moments, without fatigue to the patient, and the therapeutic properties generally are those of apomorphine, but in less degree (Dujardin-Beaumetz). The *hydrochlorate* of aponarceine is an emetic, but not in use.

PARALDEHYDUM—PARALDEHYD.

Paraldehyd is a polymeric modification of aldehyd; is soluble in water (1 in 10), less soluble in hot water than in cold; is a hypnotic in doses three times as large as of chloral. According to M. Peretti it exercises no injurious action on the heart, even in large doses; the hypnotic effect is produced rapidly, with-

CODEINE.—This more recently is usually employed in the form of the hydrochloride, melanco-

B. H. ~~_____~~

 Ten ~~_____~~

Ph water and be added to "Codeine"

and the fact that the American people are not aware of the extent of the problem. The American people are not aware of the extent of the problem. The American people are not aware of the extent of the problem.

powder of bitter taste; is sparingly soluble in water, but dissolves in ether, alcohol, or chloroform. It undergoes changes at very high temperatures—100° C. and upwards. The PEREIRINE of commerce is an impure product; of its salts, the *hydrochlorate*, *sulphate*, and *valerianate* are the only ones which have as yet been used therapeutically.

For hypodermic purposes the neutral hydrochlorate alone is capable of being used with advantage; it crystallizes in quadrangular prisms of a vermillion tint, soluble in all proportions in water. Like all pereirine salts it has an intensely disagreeable bitter taste, but no local irritant action; it abolishes the functional activity of the central grey nervous substance, and more especially of the grey matter of the medulla and spinal cord (abolition of voluntary movements, afterwards of the reflexes, and finally of the excitability of the motor and sensory nerves).

[One of the numerous substitutes for quinine, it is almost unknown in Great Britain and the United States, but in the West Indies is possessed of no small popularity as a febrifuge and antispasmodic, and employed hypodermically as well as *per orem*.—ED.]

In medicinal doses it gives rise neither to intellectual nor sensorial troubles, its influence on the circulation being trifling and irregular.

B Hydrochlorate of pereirine, 1 to 2 grains,
Camphor water, up to 30 minims.

Dose, 15 minims in each arm.

of respiratory movements, gastric disorder (nausea, vomiting) noises in the ears, sometimes feeling of pressure on the nucha. In *poisonous* doses, frequent respirations, finally loss of reflex excitability, total paralysis, and death—frequently with hyperæmia and œdema of lungs. These salts are antiseptic, and among the many cheap substitutes for quinine.

The tartrate, *hydrochlorate* and *citrate* of quinolin are the only salts used for hypodermic injections, and are painful, and often followed by extensive infiltration of the cellular tissue. The tartrate is preferable for this purpose; is of a dull white color, has a distinctive odor, and a taste analagous to that of bitter almonds; is soluble in alcohol, and insoluble in ether. The medium dose hypodermically for an adult is from 3 to 16 grains. Quinolin salts have been used with varying and uncertain results in intermittent fever, neuralgia, typhoid fever, whooping cough, tuberculosis, pneumonia, erysipelas, and septicæmia. Currie frequently used tartrate of quinolin, by internal administration, as an antipyretic in pneumonia and erysipelas. Its effects on the pulse, as tested by sphygmographic tracings, and on the temperature, closely resemble those which follow on the administration of kairin.

KAIRIN is a yellowish-white synthetic crystalline derivative of quinolin, very soluble in water and alcohol, slightly so in glycerin, insoluble in ether; it is a *hydrochlorate of oxy-ethyl-quinolin*. An anti-putres-

cent, it slows the pulse and respirations, lowers arterial tension, and diminishes temperature by vaso-motor dilatation. There is a diminution of general sensibility and motility in proportion to the amount of the dose; convulsions, contractures, pupillary contraction and paresis, occur only after large doses. Profuse sweats coincide with fall of temperature, and rigors with its rise; there may be cyanosis, hæmaturia, and diminished excretion of urea. The respiratory capacity of the blood is diminished with reduction of hæmaglobin to methæmaglobin. Elimination is effected by the kidneys with dark-green discoloration of the urine.

M. Querole used hypodermically in doses of $1\frac{1}{8}$ to 8 grains in 14 minims of water. These injections did not give rise either to local or general ill-effects. The same author states that at the ordinary temperature, kairin will only dissolve to the extent of $1\frac{1}{2}$ grains in 16 minims of water. Stronger solutions can be prepared by gradually raising the temperature of the water, which will remain clear at a temperature of 95° F. Deep injection is necessary. Kairin has been mostly used as an antipyretic, but the unpleasant effects, such as profuse perspiration and rigors, to which it gives rise, and the very temporary influence it exerts on the temperature, render it less useful than the cinchona salts.

ANALGESIN. A powerful antipyretic derived from quinolin, with properties identical with kairin, though

its action is more prolonged and intense. Also an antiputrescent and hæmostatic. It arrests fermentation in yeast, and germination. It has been claimed, beside a moderate amount of perspiration, vomiting, and pharyngeal constriction, that it produces no markedly objectionable effects; that collapse never arises from its administration. [This last, however, has recently been unequivocally proven false, and we now know that the drug is apt to prove dangerous even in moderate doses. Being a *patented* medicine, advertised in the secular press as a panacea, the less the medical profession have to do with "*antipyrin*," the better.*—ED.]

Used by a great many observers, it has chiefly been employed in the treatment of the symptom "fever." Ranke was the first to use it hypodermically. [Made in the gluteal region, it has been claimed that injections cause only transient pain without any inflammatory reaction, but Erb, Hoffer, and Huchard gave up the use of hypodermic injections on account of the intense suffering induced.†—ED.]

**Analgesin* is unpatented antipyrin, and identical in every way with the latter, at half the price. So, too, *Acetanilid* is unpatented antifebrin.—ED.

[† Dr. L. Bach, from a study of one hundred cases of various diseases of a neuralgic character, says:

1. Subcutaneous injection of analgesin as regards the production of local pain, is of the most varied action, the result evidently depending upon the individual disposition of the

THALLIN also is derived from quinolin. All the salts readily dissolve in water, but the *hydrochlorate* readily undergoes change by exposure to the light. The *sulphate*, which is generally employed, dissolves in five times, and the *tartrate* in ten times, its weight of cold water. *Ethylthallin* and its salts are equally soluble in water.

Jaksch employed these salts (and the *hydrochlorate* of ethylthallin in doses of $3\frac{1}{2}$ to $10\frac{1}{2}$ grains) as antipyretics. The fall of temperature was considerable, amounting to several degrees, but sweats frequently occurred coincidentally therewith, succeeded by rigors and an ultimate rise in temperature; also have been observed increased blood pressure and fall of

patient. 2. No difference between the first and subsequent injections can be made out. 3. Whenever possible, the injection should be made into the tissue of the muscles, since in this locality it is, at any rate, no more painful than in the subcutaneous tissue, and infiltration is avoided. 4. Previous injections of cocaine are to be recommended. 5. As recommended by Liebreich at the last Wiesbaden Congress, the injection should be made as near as possible to the seat of pain. An apparent exception to this rule is found in the fact that in hemicrania and orbital neuralgia, relief will frequently follow with astonishing rapidity the use of injections into the deltoid muscle. 6. There is no probability of analgesin ever being regarded as a substitute for morphine. 7. Injections of analgesin arrest pain in hemicrania and muscular pain, especially in lumbago and neuralgia of the sciatic and trigeminal nerves. In most cases the relief is permanent; in others it lasts six to eight hours. 8. In articular rheumatism it seems to be almost

pulse and respiration; collapse never occurred. The minimum of temperature was reached in from two to three hours after exhibition of the drug, which, further than this, exercised no influence on the progress and duration of the various diseases for which it was prescribed.

The effects produced by thallin last longer than those obtained from kairin, and it is more active than analgesin, and the dose is smaller and rarely induces vomiting ($3\frac{1}{2}$ grains of thallin = 16 grains analgesin). In doses of seven grains, thallin does not produce any symptoms in a healthy man, and all its salts arrest fermentation. Elimination is by the kidneys. Messrs.

a specific, as, in the single case in which it failed to give relief, failure can be regarded as attributable to the complications existing in the case. 9. No difference is evident as regards its influence upon acute or chronic pain; it appears to act equally well in both cases. 10. Five cases in which chills, cold sweat, palpitation of the heart, and symptoms of syncope followed its use, show that its employment should be carefully watched, although in no cases were the symptoms severe enough to cause any anxiety. Since in none of these cases was cocaine injected with the analgesin, the results are attributable to the latter alone. 11. Only in the rarest instances will these injections fail to produce some improvement.

In most cases Dr. Bach employed a solution of analgesin made in boiling distilled water. In other cases he employed a solution consisting of 150 grains each of analgesin and water, in which 3 grains of cocaine were dissolved, the latter solution being ordinarily less painful in its employment. —ED.]

Brouardel and Loye have demonstrated it has the same *modus operandi* as kairin, and acts, namely, by destruction of the hæmoglobin.

M. Grocco, of Pavia, used *acetate* of thallin, which he found was a powerful antipyretic in doses of 4 grains. The symptoms observed after its use were: Profuse sweats, and diminution in the frequency of the pulse and respirations. In some cases where there was intolerance of the drug, he administered without inconvenience by the hypodermic method in doses of $1\frac{1}{2}$ grains. The average duration of the action of this salt varied between four and six hours. M. M. Mingazzini and Pisenti used the *sulphate* and *tartrate* in water hypodermically, the latter declaring them "perfectly safe."

℞ Sulphate of thallin, 3 grains.

Distilled water, up to 16 minims.

As there is a precipitate from this solution in the cold, it is necessary always to use it warm; it does not cause abscess or other local accident. In a dose of $1\frac{1}{2}$ grains, which is equivalent to 4 grains administered *per os oris*, the lowering of the temperature reached 2.1° to 2.3° , C. and persisted for a period of six to nine hours.

"Prof. Jaccoud considers thallin the most valuable antipyretic we possess, and states that by its systematic use in small doses, it is possible to maintain a permanent condition of apyrexia in febrile cases."—(Currie.)

pulse and respiration; cooling of the body to a minimum of temperature three hours after exhibition. It is a derivative very sparingly used, more so in alcohol and water than in resorcin and causes rapid

The effects produced by 3 grains securing an immediate effect. Doses of 6 to 9 grains are given with greater certainty, and at intervals of 3½ grains of the pulse. In doses of seven grains symptoms of excitement are not observed. Its ready solubility in water makes it suitable for hypodermic use, never causing more pain than

a specific, as, in the single use, failure can be regarded as a failure existing in the case. 9. Dose, 3 grains, influence upon acute or chronic cases, up to 112 minims. well in both cases. 10. Dose, 3 grains of the salt. palpitation of the heart, as the use of hydroquinon reduces use, show that its employment is without any unpleasant although in no cases were it cause any anxiety. Since it is found that 5 to 10 minims of a solution could be injected into each tibia. 11. Only in cases of high fever, tions fail to produce some effect from repeated doses,

In most cases Dr. [Name] kept down by it continuously. The statement is open to question, as a solution consisting of 3 grains in water and water, in which 3 grains could be taken, the latter solution being ordinary.

—ED.]

RESORCINUM. RESORCIN.

isomeric with hydroquinon and *pyrocatechin*; dissolves to almost any extent in water, and is also soluble in ether, alcohol, glycerin, and vaselin; is insoluble in chloroform and carbon sulphide. Aqueous solutions when exposed to air and light soon acquire a brownish color, without appreciable alteration, according to the degree of concentration. It possesses practically the same properties as carbolic and salicylic acids, but its toxic effects are less than those of the former.

Resorcin brought in contact with albuminous substances, causes immediate coagulation forming an *albuminate* of resorcin. Slightly concentrated solutions are possessed of caustic properties, causing opacity, with white discoloration of the cellular and muscular tissues; it sometimes also produces noises in the ears and vertigo. Elimination is chiefly by the kidneys, and rapidly performed.

R Resorcin, 5 to 20 grains.

Distilled water, up to 112 minims.

Dose, 10 minims ($\frac{1}{2}$ to 3 grains), which do not give rise to local trouble.

In inveterate sciatica, intramuscular injections in a 10 or 20 per cent. solution, often produce healthy local derivation. A 5 per cent. solution has been used hypodermically by M. M. Bogusch, Stibnewski, Ugo Basse, and Cattani, without ill results, and with claimed definite and rapid effects.

"Dr. Murrell found that a dose of \mathcal{A} ij of resorcin administered in one dose to a young woman who was suffering from asthma, produced alarmingly toxic phenomena. She was, in a few minutes, in a profuse perspiration from head to foot; there was complete loss of voluntary power and reflex action, the pulse at the radials was weak and thready, and the temperature in the axilla only 94° F. Restoratives were applied, consciousness was soon restored, and the temperature gradually returned to the normal. Dr. Andrew first used hot baths as restoratives, but latterly has had recourse to the vintages of Bordeaux and Burgundy, and finds that under the influence of these stimulants the toxic symptoms speedily vanish. Dujardin-Beaumetz, who was the first in France to use resorcin, has entirely discontinued its internal administration; he says that he was led to abandon its use in rheumatism and typhoid, not only on account of its powerlessness, but also on account of the toxic phenomena to which it gave rise. 'While recognizing the fact that resorcin is not so poisonous as carbolic acid, I came to the conclusion that it was a dangerous remedy as an antipyretic, for I observed in patients suffering from typhoid fever who were treated thereby, the same depression of the vital powers, the same adynamia, and the same pulmonary congestion, which I had previously seen in patients who had been treated by carbolic acid.'" (Currie).

SCILLIAINUM AND SCILLIPICRINUM.

SCILLIAIN OR SCILLITOXIN AND SCILLIPICRIN.

Glucosides from squill, the first with the same action as digitalin when injected hypodermically; the second has been employed in the same way as a diuretic.

℞ Scillipicrin, 1 grain.

Distilled water, up to 12 or 60 minims.

Dose, $\frac{1}{3}$ to $\frac{1}{2}$ grain of the active principle.

SODIUM SALTS.

The hypodermic injection of saline solutions causes sharp suffering followed by induration, and sometimes even by abscess, though Bertin and Gray state that when preceded by an injection of atropine sulphate, common salt does not cause pain.

Injections of SODIUM CHLORIDE are sometime administered in the treatment of neuralgia, sciatica, lumbago, angina pectoris, torticollis, and muscular spasms. M. Lubanski has administered in the diarrhoea of phthisis to aid appetite and digestion, and never saw local accidents result. A 1 to 20 solution has been used for hypodermic transfusions. [Has been used in cholera, and Bright's disease, also with claimed satisfactory results.—Ed.]

IODIDE OF SODA is preferable to potassium iodide for hypodermic use, the dose being about one-third that administered by the mouth.

"Dr. Murrell found that a dose of resorcin administered in one dose to a patient who was suffering from asthma, produced the same phenomena. She was, in a few minutes, covered with perspiration from head to foot, and causes intense loss of voluntary power and the radials was weak and the patient was unable to arrest fermentation in the axilla only 94° F. but Kolbe and applied, consciousness was restored, and the temperature gradually returned to normal. The patient drew first used hot water, but resorcin causes copious perspiration, and has had recourse to it, and tinnitus aurium. Burgundy, and for convulsions and asphyxia these stimulants are contra-indicated. Elimination Dujardin-Beaume must not be forgotten resorcin, has been contra-indicated in alcoholism; he has also cardiac adynamia and renal in rheumatism.

powerlessness. Domingos and Freiré, with phenomena of asphyxia, local troubles are very the fact that the solution is recently diluted; hence, when it acid, the dose must be done remedy. The punctures—M. Collard had to suffering from the local injury to which by the authors recommend in yellow solution of 24 grains and upwards in the yellow solution of 24 grains in the second stage.

Take of sodium, 4 grains.
 Dissolve in water, up to 15 minims.

Salicylate of sodium should be perfectly neutral, and the solution should be made (and filtered) only as required. Caffeine is sometimes associated with benefit.

Gubler used GLAUBER'S SALT hypodermically, but without producing purgation. Luton used $1\frac{1}{2}$ grains in 16 minims of water, with slight laxative effect.

SOLANINA.—SOLANINE.

The therapeutic properties of this alkaloid have only recently been studied. It is obtained from potato shoots, and also from the parings of very young or very old potatoes. Solanine belongs to the same class of poisons as atropine, eserine, and pilocarpine, and forms salts with acids. The *hydrochlorate*, which is a gelatinous body readily soluble in water, may be given in doses ranging from one to three or four grains daily. Dr. Genenil regards it as a narcotic to the medulla, spinal cord, and nerve-trunks, and as a paralyzant to the terminal ends of sensory and motor nerves. It does not produce cerebral congestion, but in poisonous doses is said to affect the cerebral cortex, causing headache, delirium, cyanosis, and somnolence. It is best administered hypodermically, as it is apt, when administered by the mouth, to cause nausea, vomiting, and a desire to defæcate; it does not produce the unpleasant after-effects which are apt to follow upon the administration of morphine, and it has no cumulative action.

Solanine may be prescribed to combat excitement, spasm and pain. It has produced excellent results in the treatment of sciatica and rheumatic neuralgia, and has proved of service in the treatment of gastritis, acute dyspepsia, the vomiting of pregnancy, bronchitis, bronchial asthma, cardiac asthma, and emphysema. It has been used in cystitis, prurigo, and in some nervous affections in which restlessness and insomnia were prominent symptoms. It may be safely prescribed for young children and old persons" (Currie).

SPARTEINA.—SPARTEINE.

Sparteine is an oily fluid alkaloid extracted from *Cystitis scoparis*, the shoots of which contain a non-alkaloidal principle, SCOPARIN. In physiological properties they are supposed to closely resemble coniine.

℞ Sparteine sulphate, 2 grains.
Distilled water, 112 minims.

The diuretic effects appear after doses of $\frac{1}{4}$ of a grain (16 minims of solution).

℞ Scoparin, 6 grains.
Glycerin, 30 minims.
Distilled water up to 120.
Dose, 5 minims (= $\frac{1}{4}$ grain).

STRAMONIUM.

DATURINE, chemically identical with atropine, is sparingly soluble in water (1 in 283), freely so in alcohol, less so in ether. It is antagonized by eserine.

"According to Waring, daturine may be given hypodermically (in acute mania) in doses of $\frac{1}{120}$ to $\frac{1}{60}$ of a grain, but is less useful than atropine or hyoscyamine" (Currie.) The *extract* of stramonium is usually preferred, as it possesses no local effect.

℞ Extract of stramonium (P., D. & Co. *assayed*), 1 gr.,
Camphor water, up to 80 minims.

Dose, 5 to 20 minims, $\frac{1}{8}$ to $\frac{1}{4}$ grain.

It has been employed in preliminary emphysema (Lorent), neuralgia (Oulmont and Laurent), tetanus, shaking palsy and various forms of tremor.

In small doses it produces dilatation of the pupils, dryness of throat and mouth, dysphagia, intense thirst, slight increase of pulse and respirations, increased blood pressure, increased intestinal contractions, elevation of temperature not necessarily constant, heaviness of the head, slight excitement with tendency to movement (the legs are unsteady and give way), lassitude, occasional delirium, profound sleep; also may induce visual troubles, erotic dreams, pricking sensations and tremor. A *larger* dose produces a condition analogous to drunkenness; the visual troubles are decided, the iris completely invisible (never amblyopia); also dysphagia, dryness and burning of the throat, hoarseness of the voice, sometimes absolute aphonia, rapid respiration which becomes jerky and sighing, rapid cardiac action, fall of temperature and of arterial tension, intestinal paralysis, pallor, and chilliness with clammy sweats of the extremities. Some-

Solanine may be prescribed in the treatment of spasm and pain. It has produced sleep without the treatment of sciatica and rheumatism. It has proved of service in the treatment of dyspepsia, the vomiting of pregnancy, heartbeats; perichial asthma, cardiac asthma, and in the treatment of poisonous doses. It has been used in cystitis, prurigo, and in the treatment of the great affections in which restlessness is a prominent symptom. It may be used in the treatment of the capillary prominent symptoms. It may be used in the treatment of vascular young children and old people.

SPARTEINA.

Sparteine is an oily fluid. *Scoparia feifolia*, is a *Cystitis scoparis*, the shoots of which contain the alkaloidal principle, SCOPARINE. The Italian physicians they are supposed to be tonic, and ascribe to it

R Sparteine sulphate. M. Ambrosoli used the Distilled water, (1 spirit) hypodermically, in

The diuretic effects of sparteine, in cases of constitugrain (16 minims of solution) only a very trifling local

R Scoparin, 6 gr. began with the tincture Glycerin, 30 r. of water, and then used Distilled water tincture; a dose of sixteen

Dose, 5 minims (16), but gave rise to a hard in-

ST. have used tayuya with success,

DATURINE, chem. Dr. Eulenburg, who has sparingly soluble in the tincture of M. Ubicini, has cohol, less so in etia benefit to result from its use in

srs. Geber, Concetti, Pelizarri, Mar-
i, Rasori, and Zeissl obtained similar
its, while Prof. Sigmund, of Vienna,
ayuya was not only inefficacious but posi-
erious. TAYUAYIN (the *trianospermin* of
s a drastic, purgative principle.—ED.]

PINTHINÆ OLEUM—OIL OF TURPENTINE.

pure oil of turpentine, has been administered
odermically in doses of sixteen minims by M.
ton, but without success, in a case of obstinate
iatica; it caused intense pain, and was followed
by abscess. Turpentine injections have been used
successfully in the treatment of carcinomata, sarco-
mata, and other neoplasms, by Vogt. [It should be
remembered that turpentine in large doses in damp
or cold weather tends to supply fuel for the evolution
of animal heat rather than to exert any therapeutic
action.*—ED.]

URETHAN.

An ethylic ether of carbaminic acid. "It occurs
in the form of white crystals, free from odor, and is
readily soluble in water. Von Jaksch considers it
peculiarly suited for administration to children, and
most useful in simple uncomplicated insomnia, though
apt to prove inefficient when there is acute pain.

"Although as much as 30 to 60 grains may be

* See Vaseline, page 123.

times, from the beginning, hallucinations, g
vulsions, frequent desire to urinate, with
spasm, coma, urgent thirst, and headache
In *very* large doses, the phenomena res
mittents, and there is arrest of the hear
ipheral sensibility is only blunted in poi
Daturine exerts its action especially
sympathetic; in small doses it diminish
circulation—in strong doses it deter
paralysis (Oulmont and Laurent,)

TAYUYA.

[Tayuya, the root of *Triana*
drastic purgative and emetic, exc
system, and an active depurative
cians use also as a bitter ton
almost magical properties. M
tincture (1 in 4 of proof sp
doses of six to fifteen min
tional syphilis, and observed
reaction. Prof. Gamberini
diluted with half its bulk
injections of undiluted t
minims was well borne, b
lent swelling. Many oth
ing these observers, ha
especially in scrofula.
perimented with the
not found any real b

istence, and he states
it as a vehicle, give rise
when injected hypoder-
however, that the vaselin
The best test for im-
acid, which does not blacken
and *turpentine* have thus
in phthisis, the latter in
following formulæ:

ypsol, 5 parts;
lin, 20 parts (*by weight*).
minims to 3 drachms, to be injected
part of the thigh.
lot's turpentine, 75 minims.
vaselin, 5 drachms.
in 15 minims to 2½ drachms daily.
phenol, iodine, and *camphor*, may in
injected in the strength of one per cent.;
of per cent.; and *menthol* in 10 per cent."

VERATRIINUM—VERATRIIN.

triin is insoluble in water, soluble in alcohol
Doses of $\frac{1}{14}$ to $\frac{1}{4}$ of a grain suffice to pro-
decidedly toxic phenomena.

is an antipyretic, and when administered by
mouth produces a sensation first of heat, then of
ing, nausea, vomiting, colic, diarrhœa, tingling in
limbs; the respiratory movements become infre-
quent and painful; the pulse, which in the first place

given to adults, Dr. Saundby obtained good result from the exhibition of 2-grain doses in cases of cardiac insomnia. Mr. Jackman used in 4-grain doses every 2 hours with success in a case of tetanus, after chloral, (in 20-grain doses every 3 hours), he failed to do more than relieve pain. Dr. Sprin found hypodermic injections painless even when grains were introduced, and Dr. Savage administered in cases of insanity in doses of 5 grains subcutaneously." (Currie).

VALDIVINUM. VALDIVIN.

A principle obtained from the seeds of *Valdivia*, very sparingly soluble in water (1 in 100 parts), soluble in alcohol (1 in 60 parts), less soluble in absolute alcohol (1 in 190 parts), dissolves in chloroform, but is insoluble in ether. It has pronounced poisonous properties, but its effects are slow in developing. M. Restrope, from whom we derive all our information, has noticed vomiting after a dose of $\frac{1}{17}$ of a grain, the largest dose that has been employed. Physiological researches are incomplete, and the drug requires further investigation.

R Valdivin, 1 grain
Distilled water, 525 grains.

Dose, 15 minims ($= \frac{1}{17}$ grain). Causes slight vomiting sometimes at the point of puncture.

VASELINUM. VASELIN.

"M. Meunier considers that vaselin is absorbed very rapidly through the body, its

n
to
ed
He
ex-
rita-

(iride)
ypod-
bluted;
reased.

ORIDE OF
tica. [It
ver, and is
ughing.—

tered appo-
nent of con-
is not fixed



INDEX.

A.

	PAGE.
Abortion, threatened.....	46
Abscess, opening	54
Absorption of injections.....	4
Accidents to solutions.....	5
Acetanilid.....	108
Acetic ether.....	66
Acid, benzoic.....	15
boric.....	5
carbaminic.....	121
carbolic.....	16, 123
poisoning by.....	16
gallic.....	21
hydrocyanic	18
osmic.....	20
phenic.. ..	16
phosphoric.....	21
Prussic.....	18
poisoning by.....	20
sclerotinic	65
sulpho-ichthyolic.....	82
tannic.....	21
Aconite	20, 22,
normal liquid.....	22
poisoning	24, 27
tincture	27
<i>Aconitina</i> (aconitine).....	22
sulphate	22
Acupuncture.....	1
Adynamia.....	26, 44, 85, 90

	PAGE.
Boric acid.....	5
Brandy.....	25, 44
camphorated.....	44
Bright's disease.....	46, 57, 81, 115
Bromide of eserine.....	42
of potassium.....	105
Bromidrosis.....	84, 85
Bromine.....	38
Bromo hydrate of caffeine.....	40
conicine.....	55, 56
quinine.....	52
Bronchial asthma.....	101, 118
Bronchitis.....	22, 56, 84, 118
chronic.....	15, 45, 101, 118
Broncho-pneumonia.....	100
Brucine.....	91
Buboes.....	83

C.

<i>Caffeina</i> (caffeine).....	38, 117
bromo-hydrate.....	40
citrate.....	40
hydrochlorate.....	40
sulphate.....	40
Calabar bean.....	37, 40, 41
poisoning by.....	37, 43, 89
<i>Calabarina</i> (calabarine).....	40, 42
Calomel.....	5, 73
Camphor.....	43, 123
monobromated.....	45
poisoning by.....	44
water.....	5
Camphorated brandy.....	44
Cancer.....	31, 100, 121

	PAGE.
<i>Amygdala ol. dulce</i>	44, 94
Amyl nitrite.....	27, 54, 92
poisoning by	28, 92
Anæmia.....	25, 44, 58
cerebral.....	88
Anæsthesia, general.....	33, 47, 83
hysterical.....	35
local.....	48, 83, 91
Anal fissure.....	82
Analgesinum (analgesin).....	107, 108
Anasarca, general ..	46
Angina pectoris.....	20, 22, 56, 87, 88, 92, 115
tonsillaris.....	100
Anise, ammoniated with alcohol.....	25
Analodyne, Hoffman's.....	66
Anorexia.....	59, 67
ANSTIE, DR.....	I
Anthrax.....	18
Anti-ferments.....	15, 16, 49, 66, 89, 107, 116
-phlogistics.....	82
-putrescents.....	15, 16, 49, 53, 66, 89, 106, 116
-pyretics.....	47, 50, 102, 106, 112, 114, 123
the more valuable.....	111
-septics.....	83, 93
-sialagogues..	36 59
-spasmodics.....	45, 103
-sudorifics.....	36, 50, 83, 100
-zymotics.....	49, 58
<i>Antimonium tartratum</i>	29
Antimony tartrate.....	29
Antipyrin.....	108
Aortic diseases.....	34, 88
incompetency	34
Aphrodisiacs.....	47, 67

	PAGE.
Cathartics.....	26, 117
drastic.....	121 124
Catheter fever.....	22
introduction of.....	22
<i>Cedrinum</i> (cedrin).....	47
Cephalalgia.....	22
Charbon.....	18
Cherry-laurel water.....	5, 19
poisoning by.....	20
Chinoidine.....	53
Chloral croton.....	48
hydrate.....	48, 89
poisoning by.....	89, 91 100
Chloride of gold.....	34
mercury and urea.....	73
sodium.....	70, 115
zinc.....	125
Chloroform.....	28, 48
poisoning.....	28, 89, 100
water.....	5
Cholera.....	27, 32, 39, 46, 51, 53, 59, 85, 91, 98
infantum.....	53
Chordee.....	22
Chorea.....	31, 32, 36, 42, 45, 56, 60, 79, 81
Cinchona salts.....	49
Cinchonidia sulphate.....	50
Cleaning needles.....	13
<i>Cocaina</i> (cocaine).....	53, 109, 110
hydrochlorate.....	53, 70
poisoning.....	28, 89, 91
salicylate.....	54
<i>Codeina</i> (codeine).....	100
phosphate.....	100
Coffee.....	25, 39

	PAGE.
Coffee, fluid extract of	39
<i>Colchicina</i> (colchicine)	54
<i>Colchicum seminis</i>	55
Colic.	29
Collapse	15, 27
Coma uræmic	85
Committee Report of Medico-Chirurg. Society	2
Condurango	55
<i>Coniina</i> (coniine; conicine)	42, 55
hydrobromate	55, 56
Conium	55
poisoning by	56
<i>Convallamarina</i> (convallamarine)	58
<i>Convallaria majalis</i>	57
fluid extract	57
Convulsions	53, 56, 81, 115, 118
infantile	86
epileptiform	81
<i>Cotoinum</i> (cotoin)	58
Cough	101
convulsive	56
nervous	20
of phthisis	20
whooping (see Pertussis).	
<i>Coup de soleil</i>	27, 53
Croton chloral	48
Croup	56, 73, 100
catarrhal	22
Curare	59
poisoning by	60, 89
<i>Curarina</i> (curarine; ourari)	59, 61, 89
CURRIE, DR. (Preface), 15, 21, 29, 34, 37, 38, 39, 46,	
48, 52, 54, 73, 80, 81, 85, 88, 89, 92, 93, 94, 101, 106,	
111, 112, 114, 118, 119, 122,	123

	PAGE.
.....68,	70
.....	19
.....	20
.....	118
.....15,	118

.....	118
.....	119
.....	118
.....35,	119
.....	90
.....	58
.....	81
.....79, 80,	90
.....	81
.....18,	101
.....36, 83,	100
.....83,	100
.....58, 83,	115
.....	62
.....57,	62
.....	62
.....	62
.....18, 85, 100,	105
.....	91
.....	94

~~.....~~ Nephritis, Renal,

.....31, 32, 46, 82, 83,	118
.....30, 39, 115,	118
.....	28
.....	64
.....	96

	PAGE.
Dog, effect of strychnine on.....	89
Donovan's solution.....	32
<i>Duboisina</i> (duboisine).....	63
poisoning by.....	63
sulphate.....	63
Dysmenorrhœa.....	28, 45, 75, 76, 77
Dyspepsia.....	20, 91, 118, 125
obstinate.....	125
Dysphagia.....	54
Dyspnœa.....	33, 34, 56
cardiac.....	57

E.

Eclampsia.....	20, 56, 84
Ecthyma.....	83
Eczema.....	20, 46
chronic.....	32
Emaciation.....	78
Emetics.....	29, 34, 64, 99, 100, 101, 122
<i>Emetina</i> (emetine).....	63
nitrate.....	63
poisoning by.....	64
sulphate.....	63
Emphysema, pulmonary.....	56, 91, 118, 119
Endermic medication.....	I
Endometritis.....	76, 77
Enteralgia.....	81
Enuresis.....	36, 40, 91
Epilepsy.....	28, 32, 45, 56, 60, 79, 80, 81, 84, 87, 94, 100
Epileptiform tic.....	88
<i>Ergotum</i>	65
extract (ergotin, ergotinine).....	65, 76
normal liquid.....	65
Erysipelas.....	17, 106

Cyanide of

2A

Cystitis scro-

Cystitis..

Datura .

Daturin

Death :

Debilit:

Deliri

Dem-

Dial

Dia:

Dia

Di:

Di

Di:

D

I

I

..... 1

..... 12

..... 16

..... 26, 12

..... 6

.....

.....

..... 11

..... 29, 10

..... 2

..... 3

..... 3

..... 5

..... 6

..... 7

..... 11

.....

.....

..... 7

.....

.....

..... 6

..... 6

..... 6

.....

..... III, 11

..... 9

..... 44, 85, 9

..... 2

.....

	PAGE.
Fever, inflammatory.....	22
intermittent.....	22, 47, 50, 52, 53, 83, 84, 106
puerperal.....	22, 66, 85
remittent.....	53
septic (see Septicæmia).....	
symptomatic.....	124
typhoid.....	15, 18, 22, 53, 85, 94, 106, 124
typhus.....	15, 26, 52
yellow.....	101, 116
Formiamidate of mercury.....	71
Forminate of quinine.....	52
Formyl ter-chloride,.....	48
ter-iodide.....	82
Fowler's solution.....	31
Fox-glove.....	62
Fungi poisoning.....	36

G.

Galactagogue.....	85
, anti-.....	86
Gallic acid.....	21
Gangrene.....	38, 83
Gastralgia.....	20, 81, 83
Gastric catarrh.....	55
diseases.....	20, 46, 55, 81, 91, 100, 118, 125
ulcer.....	46
Gastrodynia.....	100
<i>Geissospermum Vellonii</i>	102
Gelsemin (concentration).	68
<i>Gelsemina</i> (gelsemine alkaloid).....	68
hydrochlorate.....	68
<i>Gelsemium sempervirens</i>	67
normal liquid.....	68
poisoning by.....	67

	PAGE.
Generative desire.....	67
Genital congestion.....	76
Glandular enlargements.....	82
Glaucoma.....	26
Glauber's salt.....	117
Glonoin.....	87
poisoning by.....	87
Glycerin as a solvent.....	5
Glycocholate of mercury.....	71
Goitre.....	82
exophthalmic.....	63
Gold chloride.....	34
Gonorrhœa.....	22
Gout (see Rheumatism also).	15, 31
Gummata, mucous inter-digital.....	83

H.

Hæmoptysis.....	21
Hæmorrhage.....	36, 77
renal.....	30
intra-uterine.....	27, 46, 76
Hæmostatics.....	36
Hasisch or hashish ...	45
Hay fever (see Autumnal catarrh)	
Heart (see Cardiac).....	
Heat stroke.....	27, 53
Hemeralopia.....	91
Hemicrania.....	54, 109
Hemiplegia (see Paralysis).....	
Hemp Indian.....	45
poisoning by.....	47
Henbane.....	78
Hernia, strangulated.....	39
<i>Homatropina</i> (homatropine).....	37

	PAGE.
<i>Homatropina</i> muriate.....	37
HUNTER, DR. CHAS.....	1
Hydrargyrum and preparations.....	68-73
<i>Hydrastina</i> (hydrastinine).....	73
muriate.....	78
poisoning by.....	79
<i>Hydrastis Canadensis</i>	74
Hydrochinon	112
Hydrocyanic acid.....	18
poisoning by.....	20
Hydroquinon.....	112
<i>Hyoscina</i> (hyoscine).....	80
hydriodate, hydrobromate, hydrochlorate.....	80
<i>Hyoscyamina</i> (hyoscyamine).....	35, 78
poisoning by.....	80
Hyoscyamus.....	78
normal liquid.....	79
poisoning by.....	80
tincture.....	78
Hypnotic.....	96, 101 (2), 105, 116, 117
a reliable.....	80
Hypodermic injections.....	5
how to make.....	6
physiological activity of.....	4
point for.....	7
safety of.....	3
Stockwell's method.....	6
temperature of ..	5
when inhibited.....	7
syringes.....	7
care of	9, 13, 66
Hypopion.....	85
Hysteria.....	36, 39, 46, 56, 58, 80, 87, 100
Hystero-epilepsy.....	81, 100

	PAGE.
I.	
Ichthyol.....	82
Impetigo.....	20
Impotence.....	67
Incontinence (see Urine also).....	36
Injectons hypodermic.....	5
of air.....	7
in alcoholics.....	7
into blood vessels.....	6
in the cachectic.....	7
diabetic.....	7
massage of.....	70, 71
painless.....	70
in the purpuric.....	7
temperature of.....	5
in veins.....	6
Inoculation of remedies.....	1
Insanity.....	26, 81
Insomnia.....	39, 81, 100
alcoholic.....	46
nervous.....	46
cardiac.....	122
Intestinal atony.....	91
obstruction.....	41
Intoxication (see Alcoholism and Morphinism).....	
Introduction.....	1
Iodide of potassium.....	105
sodium.....	105
Iodine.....	123
Iodoform.....	82, 123
Ipecac.....	65
Iron preparations.....	66

J.

<i>Jaborandum</i> (jaborandi)	63, 83
normal liquid	83
Jessamine, yellow	67
normal liquid	68

K.

<i>Kairinum</i> (kairin)	106
Keratittis	26

L.

LAFARGUE, DR.	I
Laryngitis	22
Laryngismus stridulus	56, 73, 100
Laurel water	5, 19
poisoning by	20
Leiter's hypodermic syringe	7, 11
Lichen ruber	31
tropicus	20
Lilly of Valley	57
<i>Liquor ammoniæ</i>	27
<i>arsenii et hydrarg. iod.</i>	32
<i>potassii arsenit.</i>	31
<i>sodii arsenit</i>	31
Locomotor ataxia (see Tabes dorsalis)	
Lumbago	20
Lupus	31

M.

<i>Mai de mere</i>	28, 88
Mammary cancer (see Cancer)	
secretion—suppression of	86
Mania	20, 46, 53, 79, 81, 100
acute	56, 80, 119

	PAGE.
Mania, <i>a potu</i>	80, 90
hysterical (see Hysteria).....	80
puerperal	20, 80
Medico-Chirug. Society Report....	2
Melancholia.....	20, 39 54, 81
Meniere's disease	88
Meningitis.....	61
<i>Menispermum cocculus</i>	104
Menorrhagia.....	45
Menstruation irregular.....	76, 77
Mental diseases.....	20, 45, 62, 81, 101
Menthol.....	123
Mercury and salts.....	68-73
poisoning by.....	69, 84
Metrorrhagia.....	75
Migraine.....	47, 54, 88
Milk, suppression of secretion of.....	86
Monobromated camphor.....	45
Morphia alkaloid.....	95
acetate	96
and atropine.....	35
hydrochlorate.....	95, 97
meconate.....	96
muriate.....	95, 97
poisoning by.....	25, 38, 39, 96, 100
sulphate.....	96
Morphinism	38, 39 96
<i>Moschus, tinct.</i>	85
<i>Muscæ volitantes</i>	85
<i>Muscarina</i> (muscarine).....	63, 86
poisoning by.....	36, 37, 86
sulphate	86
Musk, tincture.....	85
Mydriasis.....	54

	PAGE.
Myomata.....	76
Myxœdema.....	88

N.

Nævus.....	18
<i>Napellina</i> (napelline).....	24
Nausea.....	83
Needles, care of.....	13
cleansing.....	13
obstruction of.....	13
Neoplasms.....	31, 55, 82, 100, 121
Nephritic hæmorrhage.....	30
Nephritis (see Renal also).....	30, 39, 93
acute.....	87
Bright's.....	46, 57, 81, 88, 115
chronic.....	87
delirium of.....	81
interstitial.....	88
<i>Nerium oleander</i>	94
Neuralgia.....	17, 20, 31, 32, 36, 39, 45, 53, 55, 62,
68, 87, 88, 101, 106, 108, 115, 118, 119, 124, 125	
aural.....	18
intercostal (see Rheumatism also).....	17
obstinate.....	20
orbital.....	109
rheumatic.....	118
sciatic.....	17, 28,
45, 48, 49, 55, 91, 109, 113, 115, 118, 121, 123, 125	
supra-orbital.....	17
trigeminal.....	22, 48, 109
Neuroses.....	20, 45, 46, 118
<i>Nicotina</i> (nicotine).....	86
poisoning.....	58
Night-blindness.....	91

	PAGE.
Nitrate of silver.....	30
Nitrite of amyl.....	27
Nitro-glycerin.....	87
poisoning by.....	87
Normal liquid aconite.....	22
belladonna.....	36
cannabis.....	46
digitalis.....	62
ergot.....	65
gelsemium.....	68
hyoscyamus.....	79
jaborandi.....	83
nux vomica.....	90
opium.....	95
Nutrients.....	94
Nux Vomica.....	88
normal liquid.....	90
poisoning by.....	89, 92
tincture.....	90

O.

Odontalgia.....	54, 83
Edema.....	39
malignant.....	18
pulmonary.....	93
Esophageal cancer.....	55
obstruction.....	100
Oil almond, bitter.....	19
poisoning by.....	20
sweet.....	44, 94
olive.....	94
turpentine.....	121
Oils.....	5
<i>Oleander-nerium</i>	94

	PAGE.
<i>Oleandrina</i> (oleandrine).....	94
Opium and alkaloids.....	25, 95
normal liquid.....	95
poisoning by (see Morphine).....	25, 96
Osmic acid.....	20
Ouari.....	59
<i>Ouarina</i> (ouarine).....	59
antidote to.....	60
Oxy-ethyl-quinolin muriate.....	106
Oxytoxics.....	49

P.

Packing, care of.....	8, 11
Pain (see Neuralgia, Rheumatism).....	118
acute.....	29, 121
epigastric.....	21, 81, 83
muscular.....	17, 109
rheumatoid.....	124
tabetic.....	29, 85, 91
Pancreas secretion of—to stimulate.....	59
<i>Pao-pereira</i>	102
<i>Papaverina</i> and salts.....	98
<i>Paracotoina</i> (paracotoin).....	58
<i>Paraldehydum</i> (paraldehyd).....	101
poisoning by.....	102
Paralysis.....	79
agitans.....	31, 32, 79, 119
Bell's.....	91
from compression of radical nerve.....	91
of crico-thyroid muscles.....	91
diphtheritic.....	91
hemiplegic.....	91
infantile.....	91
lead.....	91

	PAGE.
Paralysis, paraplegic.....	91
of vocal cords.....	91
writers' ("writer's cramp").....	91
Parke, Davis & Co.'s hypodermic tablets.....	
.....9, 10, 42, 52, 62, 79, 90,	99
hypodermic syringes.....	11
Pearson's sol. (<i>liq. sodii arsen.</i>).....	31
Peptonate of mercury.....	71
<i>Pereirina</i> (pereirine).....	102
Perspiration excessive.....	24, 36, 50, 83, 84, 100
foetid.....	84, 85
Pertussis.....	20, 22, 28, 45, 56, 81, 106
Phenic acid.....	16, 123
Phenol.....	16, 123
Phosphate of codeine.....	100
Phosphoric acid.....	21
Phthisis.....	18, 20, 21 (2), 22, 28, 56, 57, 81, 82, 100, 106, 123
cough of.....	20, 24, 36, 86
sweats of.....	24, 36, 37, 50, 63, 86, 104
diarrhœa of.....	115
insomnia of.....	46
<i>Physostigmina</i> (physostigmine).....	40
poisoning by.....	27, 43
<i>Picrotoxinum</i> (picrotoxin).....	104
<i>Pilocarpina</i> (pilocarpine and salts).....	83
poisoning by.....	37, 85
<i>Pilocarpine pinmatus</i> (jaborandi).....	83
Pleurisy.....	39, 56, 84
Pleuro-pneumonia.....	18
Pneumonia.....	15, 18, 56, 84, 100, 106, 124
acute.....	93
broncho-.....	100
delirium of.....	81
Poisoning by aconite.....	20, 24, 27

	PAGE.
Poisoning by aloin	26
amyl nitrite.....	28, 92
apocodeine.....	34
arsenic	32
atropine.....	37, 43, 85, 86, 91
belladonna.....	36
bitter almond oil	20
Calabar bean.....	37, 43, 89
camphor.....	44
carbolic acid	15
cherry laurel water.....	20
chloral	89, 91, 100
chloroform....	28, 89, 100
cocaine.....	28, 89, 91
curare.....	60, 89
cyanide of potassium.....	20
daturine.....	85, 119
digitalis	62
duboisine.....	63
emetine.....	64
eucalyptol.....	66
gelsemium.....	67
hydrastinine.....	74
hyoscyamus.....	80
mercurials.....	69, 84
morphine	25, 38, 39, 96, 100
muscarine.....	26, 37, 86
nicotine	58
nitro-glycerin	87
nux vomica.....	89, 92
opium	25, 96
paraldehyd.....	102
physostigmine	37, 43
pilocarpine.....	37, 85

	PAGE.
Q.	
Quebrachin and salts.....	33
Quebracho.....	33, 34
Quinidine.....	49
Quinine.....	4
amorphous.....	53
bi-sulphate.....	52
carbamide-hydrochlorate.....	52
carbolate.....	51
formiate.....	52
hydrochlorate.....	50, 51
hydrobromate.....	52
lactate.....	52
poisoning by.....	49
quinate.....	52
sulphate.....	50, 51
sulpho-vinate.....	52
tannate.....	52
valerianate.....	52
Quinodina.....	53
Quinoline and derivatives.....	105
tartrate.....	106
citrate.....	106
poisoning by.....	106
Quinon.....	112

R.

Rectal prolapse.....	90
Renal disease (see Nephritis).....	
hæmorrhage.....	30
Report of Commission of Royal Medico-Chirurgical Society.....	2
<i>Resorcinum</i> (resorcin).....	113
albuminate of.....	113

	PAGE.
<i>Resorcinum</i> , poisoning by.....	113, 114
Retina detached.....	85
Rheumatism.....	17, 20, 31, 45, 79, 115, 124
acute.....	22, 23, 53, 82, 105
articular.....	18, 53, 55, 84
chronic.....	22, 53, 82
articular.....	53, 55
intercostal.....	17
lumbago.....	20, 77, 115
neuralgic.....	118
sciatic (see Sciatica).	
Rheumatoid tetanus.....	83
Rickets.....	82
Royal Medico-Chirurg. Soc. Com. Report.....	2
Rupia.....	83
RUPPANNER, DR.....	2

S.

Salicylate of soda.....	39,	116
Salts of sodium.....		116
Glauber's		116
Sarcoma diffuse.....		31
Sciatica.....	17, 28, 45, 48, 49, 55, 91, 109,	113
	115, 118, 121, 123,	125
Scillitoxinum (scillitoxin).....		115
Sclero-choroiditis posterior.....		85
Sclerosis.....		99
Sclerotinic acid.....		65
<i>Scillainum</i> (scillain).....		115
<i>Scillipicrinum</i> (scillipicrin).....		115
Scrofula.....	82,	120
Scoparine.....		118
Sea-sickness.....	28,	80
Septicæmia.....	22, 53,	106

	PAGE.
Septic fever.....	22, 53, 106
Serous collections.....	82
Sheen's hypodermic syringe.....	9
Sialagogues.....	49, 59
Silver nitrate.....	30
<i>Simaba cedron</i>	47
<i>valdiva</i>	122
Skin diseases.....	17, 20, 31, 32, 46, 82, 83, 118
Soda salts.....	65
Society, Royal Med.-Chirurg Report.....	2
Sodium chloride.....	70, 115
iodide.....	115
sulphide.....	117
lactate.....	116
salicylate.....	116
poisoning by.....	116
<i>Solanina</i> (solanine).....	117
poisoning by.....	117
Solutions.....	5
accidents to.....	5
acid.....	4
concentrated.....	4
fungi in.....	5
neutral.....	4
preserving.....	5
Solvents.....	5
<i>Sparteina</i> (sparteine).....	118
Spasms (see Convulsions also).....	53, 56, 81, 86, 115, 118
muscular.....	115
Spirit, turpentine.....	121, 123
<i>vini recti</i>	5, 25, 44
Spleen, hypertrophy of.....	53
Stimulants.....	59
diffusible.....	25, 44, 65

	PAGE.
Stockwell's method.....	6
Stomach (see Gastric also).	
pain	20, 81, 83
catarrh.....	55
Stomachics	59
Stramonium	118
extract.....	119
poisoning by.....	85, 120
Strychnia.....	87
arseniate	93
poisoning.....	42, 44, 61, 86, 88, 89, 91, 92, 101, 102
<i>Strychnos nux vomica</i>	89
<i>toxifera</i>	59
Sulpho-ichthyolic acid.....	82
Sunstroke.....	27, 53
Sutjee.....	45
Sweats (see Antisudorifics).	
night (see Phthisis).	
unilateral.....	84
Syncope (see Cardiac diseases).....	27, 88
Syphilis.....	35, 71, 72, 73, 82, 85, 120
Syringe, care of.....	13
filling.....	11
joints.....	9, 66
Leiter's.....	7, 11
Parke, Davis & Co.'s.....	7, 11
Sheen's.....	9

T.

Tabes dorsalis.....	29, 31, 79, 85, 91
pain of.....	29, 85, 91
Tablets, hypodermic.....	9, 10, 42, 52, 62, 79, 90, 99
Tannic acid.....	21
Tannin.....	21

	PAGE.
Tartar emetic.....	29
Tartrate of antimony.....	29
Tayuya.....	120
<i>Tayuyinum</i> (tayuyin).....	121
Ter-chloride of formyl.....	48
Ter-iodide of formyl.....	82
<i>Terebinthina oleum</i>	121, 123
Tetanus.....	17, 22, 36, 45, 49, 56, 60, 85, 86, 119, 122
rheumatoid.....	83
<i>Thallinum</i> (thallin).....	111
acetate.....	109
hydrochlorate.....	109
sulphate.....	109
tartrate.....	109
Thymol.....	123
<i>Tic</i> , convulsive.....	61
<i>douloureux</i>	45, 50
epileptiform.....	88
Tincture aconite.....	27
belladonna.....	25, 36, 37
cannabis.....	45
digitalis.....	62
hyoscyamus.....	78
Indian hemp.....	45
musk.....	85
nux vomica.....	90
tayuya.....	120
Tobacco heart.....	58
poisoning.....	68
Tonics.....	47, 66, 102, 103, 120
Tonsillitis.....	22, 100
Toothache (see Odontalgia).	
Torticollis.....	115
Tremor (see also Paralysis).....	31, 32, 79

	PAGE.
Tremor, alcoholic (see also Delirium tremens)	79
senile (see Paralysis agitans).....	79
<i>Trianosperma fefolia</i>	120
<i>tinctura</i>	120
<i>Trianosperminum</i> (trianospermin).....	121
Trinitrin.....	87
Trismus neonatorum.....	42
Tuberculosis (see also Phthisis).....	18, 21, 53, 106
insomnia of.....	46
Tumors (see also Neoplasms).....	20, 119
painless removal of.	54
Turpentine.....	121, 123
Typhoid (see Fever).	
Typhus (see Fever).	

U.

Ulcers, cancerous	82, 83
phagadenic.....	83
sluggish	83
stomach.....	46
varicose	83
venereal	82, 83
Uræmia	15
Uræmic coma.....	85
Urine, incontinence of	36, 42, 91
Urari (curare).....	59, 89
Urea chloride with mercury.....	73
with quinine.....	52
Urethan	89, 121
poisoning by	89
Urticaria.....	20
Uterine atony.....	46
cancer	46, 82
contraction.....	85

	PAGE.
Uterine inertia.....	46
Uva ursi.....	39

V.

Vaginismus.....	36
<i>Valdivinum</i> (valdivin).....	122
Valerianate of ammonia.....	27
quinine.....	52
Valleix's method.....	2
Vascular tension.....	88
Vaselin.....	122
test for.....	123
Venom wounds.....	27, 47, 105
<i>Veratriinum</i> (veratriin).....	122
<i>Veratrina</i> (veratrine).....	124
<i>Veratrum album</i>	124
<i>sabadilla</i>	124
<i>viride</i>	123
normal liquid.....	125
Vertigo.....	88
Vesical catarrh.....	30
inflammation.....	30
Vomiting.....	83
functional.....	20
in pregnancy.....	36, 118

W.

Water.....	29
camphor.....	5
chloroform.....	5
cinnamon.....	97
distilled.....	5
eucalyptus.....	5
laurel.....	5, 19





LANE MEDICAL LIBRARY

To avoid fine, this book should be returned on
or before the date last stamped below.

JUN 17 1994
JUN 11 1994

U169 Bourneville, D.M.
B77 Hypodermic medication
1890 10987

[illegible]

